




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|--|---|
| (51) International Patent Classification ⁶ : C12N 15/00, 15/12, 15/62, 15/63, 15/67, 15/85, 5/10, 1/15, 1/19, C07K 14/245, A01K 61/027 | | A1 | (11) International Publication Number: WO 96/01313 |
| | | | (43) International Publication Date: 18 January 1996 (18.01.96) |
| (21) International Application Number: PCT/US95/08179 | | Manfred [DE/US]; 978 Arlington Boulevard #B, El Cerrito, CA 94530 (US). | |
| (22) International Filing Date: 29 June 1995 (29.06.95) | | (74) Agents: DECONTI, Giulio, A., Jr. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US). | |
| (30) Priority Data: | | (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). | |
| 08/270,637 1 July 1994 (01.07.94) US 08/275,876 15 July 1994 (15.07.94) US 08/383,754 3 February 1995 (03.02.95) US 08/846,814 7 June 1995 (07.06.95) US | | | |
| (60) Parent Applications or Grants | | Published | |
| (63) Related by Continuation | | With international search report. | |
| US 08/270,637 (CIP) Filed on 1 July 1994 (01.07.94) US 08/275,876 (CIP) Filed on 15 July 1994 (15.07.94) US 08/383,754 (CIP) Filed on 3 February 1995 (03.02.95) US 08/486,814 (CIP) Filed on 7 June 1995 (07.06.95) | | Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. | |
| (71)(72) Applicants and Inventors: BUJARD, Hermann [DE/DE]; Remler Strasse 9, D-69120 Heidelberg (DE). GOSSEN, | | | |

(54) Title: TETRACYCLINE-REGULATED TRANSCRIPTIONAL MODULATORS



(57) Abstract

Nucleic acid molecules and proteins useful for regulating the expression of genes in eukaryotic cells and organisms are disclosed. The invention provides a transcriptional activator fusion protein which binds to tet operator sequences and stimulates transcription of a tet operator-linked gene in the presence, but not the absence, of tetracycline (or analogue thereof). The invention further provides transcriptional inhibitor fusion proteins which inhibit transcription of a tet operator-linked gene in a regulated manner. In one embodiment, the inhibitor fusion protein binds to tet operator sequences in the absence, but not the presence, of tetracycline (or analogue). In another embodiment, the inhibitor fusion protein binds to tet operator sequences in the presence, but not the absence, of tetracycline (or analogue). The transcriptional activator and inhibitor fusion proteins of the invention can be used in combination to regulate expression of one or multiple tet operator-linked genes. Novel tet operator-containing transcription units which allow for coordinate or independent tetracycline-regulated expression of two or more genes by the transcriptional modulators of the invention are also disclosed. Kits including the components of the regulatory system are also encompassed by the invention.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria | GB | United Kingdom | MR | Mauritania |
| AU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | IT | Italy | PL | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgyzstan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SI | Slovenia |
| CI | Côte d'Ivoire | LI | Liechtenstein | SK | Slovakia |
| CM | Cameroon | LK | Sri Lanka | SN | Senegal |
| CN | China | LU | Luxembourg | TD | Chad |
| CS | Czechoslovakia | LV | Latvia | TG | Togo |
| CZ | Czech Republic | MC | Monaco | TJ | Tajikistan |
| DE | Germany | MD | Republic of Moldova | TT | Trinidad and Tobago |
| DK | Denmark | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | US | United States of America |
| FI | Finland | MN | Mongolia | UZ | Uzbekistan |
| FR | France | | | VN | Viet Nam |
| GA | Gabon | | | | |

TETRACYCLINE-REGULATED TRANSCRIPTIONAL MODULATORS

Background of the Invention

Functional analysis of cellular proteins is greatly facilitated through changes in the expression level of the corresponding gene for subsequent analysis of the accompanying phenotype. For this approach, an inducible expression system controlled by an external stimulus is desirable. Ideally such a system would not only mediate an "on/off" status for gene expression but would also permit limited expression of a gene at a defined level.

Attempts to control gene activity have been made using various inducible eukaryotic promoters, such as those responsive to heavy metal ions (Mayo et al. (1982) *Cell* 29:99-108; Brinster et al. (1982) *Nature* 296:39-42; Searle et al. (1985) *Mol. Cell. Biol.* 5:1480-1489), heat shock (Nouer et al. (1991) in *Heat Shock Response*, e.d. Nouer, L., CRC, Boca Raton, FL, pp167-220) or hormones (Lee et al. (1981) *Nature* 294:228-232; Hynes et al. (1981) *Proc. Natl. Acad. Sci. USA* 78:2038-2042; Klock et al. (1987) *Nature* 329:734-736; Israel & Kaufman (1989) *Nucl. Acids Res.* 17:2589-2604). However, these systems have generally suffered from one or both of the following problems: (1) the inducer (e.g, heavy metal ions, heat shock or steroid hormones) evokes pleiotropic effects, which can complicate analyses, and (2) many promoter systems exhibit high levels of basal activity in the non-induced state, which prevents shut-off the regulated gene and results in modest induction factors.

An approach to circumventing these limitations is to introduce regulatory elements from evolutionarily distant species such as *E.coli* into higher eukaryotic cells with the anticipation that effectors which modulate such regulatory circuits will be inert to eukaryotic cellular physiology and, consequently, will not elicit pleiotropic effects in eukaryotic cells. For example, the Lac repressor (lacR)/operator/inducer system of *E. coli* functions in eukaryotic cells and has been used to regulate gene expression by three different approaches: (1) prevention of transcription initiation by properly placed *lac* operators at promoter sites (Hu & Davidson (1987) *Cell* 48:555-566; Brown et al. (1987) *Cell* 49:603-612; Figge et al. (1988) *Cell* 52:713-722; Fuerst et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:2549-2553; Deuschle et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5400-5405); (2) blockage of transcribing RNA polymerase II during elongation by a LacR/operator complex (Deuschle et al. (1990) *Science* 248:480-483); and (3) activation of a promoter responsive to a fusion between LacR and the activation domain of herpes simplex virus (HSV) virion protein 16 (VP16) (Labow et al. (1990) *Mol. Cell. Biol.* 10:3343-3356; Baim et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:5072-5076).

In one version of the Lac system, expression of *lac* operator-linked sequences is constitutively activated by a LacR-VP16 fusion protein and is turned off in the presence of isopropyl- β -D-thiogalactopyranoside (IPTG) (Labow et al. (1990), cited *supra*). In another version of the system, a lacR-VP16 variant is used which binds to *lac* operators in the presence of IPTG, which can be enhanced by increasing the temperature of the cells (Baim et

al.(1991), cited *supra*). The utility of these *lac* systems in eukaryotic cells is limited, in part, because IPTG acts slowly and inefficiently in eukaryotic cells and must be used at concentrations which approach cytotoxic levels. Alternatively, use of a temperature shift to induce gene expression is likely to elicit pleiotropic effects in the cells. Thus, there is a need
5 for a more efficient inducible regulatory system which exhibits rapid and high level induction of gene expression and in which the inducer is tolerated by eukaryotic cells without cytotoxicity or pleiotropic effects.

Components of the tetracycline (Tc) resistance system of *E. coli* have also been found to function in eukaryotic cells and have been used to regulate gene expression. For example,
10 the Tet repressor (TetR), which binds to *tet* operator sequences in the absence of tetracycline and represses gene transcription, has been expressed in plant cells at sufficiently high concentrations to repress transcription from a promoter containing *tet* operator sequences (Gatz, C. et al. (1992) *Plant J.* 2:397-404). However, very high intracellular concentrations of TetR are necessary to keep gene expression down-regulated in cells, which may not be
15 achievable in many situations, thus leading to "leakiness" in the system.

In other studies, TetR has been fused to the activation domain of VP16 to create a tetracycline-controlled transcriptional activator (tTA) (Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551). The tTA fusion protein is regulated by tetracycline in the same manner as TetR, i.e., tTA binds to *tet* operator sequences in the
20 absence of tetracycline but not in the presence of tetracycline. Thus, in this system, in the continuous presence of Tc, gene expression is kept off, and to induce transcription, Tc is removed.

Summary of the Invention

25 One aspect of the invention pertains to an inducible regulatory system which utilizes components of the Tet repressor/operator/inducer system of prokaryotes to stimulate gene expression in eukaryotic cells. In the system of the invention, transcription of a *tet* operator-linked nucleotide sequence is kept silent in the absence of tetracycline but can be rapidly and strongly induced in the presence of tetracycline (or analogue thereof). Transcription is
30 induced by a fusion protein composed of at least two polypeptides, a first polypeptide which binds to *tet* operator sequences in the presence of tetracycline or a tetracycline analogue and a second polypeptide which directly or indirectly activates transcription in eukaryotic cells. In a preferred embodiment, the first polypeptide of the fusion protein is a mutated Tet repressor which is regulated by Tc in a reverse manner from the wild type repressor, i.e., it binds to *tet*
35 operator sequences in the presence rather than in the absence of Tc. Thus, in the absence of the inducing agent (Tc or a Tc analogue), transcription of a *tet* operator-linked nucleotide sequence remains uninduced. In the presence of the inducing agent, transcription of the *tet* operator-linked nucleotide sequence is stimulated by the transactivator fusion protein of the invention.

The inducible regulatory system of the invention has the advantageous properties that induction of gene expression is rapid, efficient and strong (e.g., typically between 1000 to 2000-fold; up to a 20,000-fold increase in expression has been observed) and the inducing agent does not have to be continuously present. Moreover, the inducing agent does not cause pleiotropic effects or cytotoxicity in eukaryotic cells. The inducible regulatory system of the invention can be applied to regulation of gene expression in cells either *in vitro* or *in vivo*, and may be particularly useful for gene therapy applications and for expression of gene products in transgenic and homologous recombinant organisms (e.g., animals and plants).

The inducible regulatory system of the invention involves at least two components: a tetracycline-inducible transcriptional activator and a target transcription unit to be regulated. Accordingly, one aspect of the invention pertains to a Tc-inducible transcriptional activator ("transactivator") fusion protein, and nucleic acid (e.g., DNA) encoding the fusion protein. A preferred fusion protein comprises a Tn10-encoded Tet repressor that is mutated at at least one amino acid position selected from amino acid positions 71, 95, 101 and/or 102, operatively linked to a herpes simplex virus virion protein 16 (VP16) activation domain (the nucleotide and amino acid sequences of such a fusion protein are shown in SEQ ID NO: 1 and 2, respectively). In one embodiment, the activation domain of the fusion protein includes about 127 C-terminal amino acids of VP16 (e.g., the fusion protein of SEQ ID NO: 2). In another embodiment, the activation domain includes at least one copy of about 11 C-terminal amino acids of VP16 (e.g., the amino acid sequence of which is shown in SEQ ID NO: 4). Other mutated Tet repressors and transcriptional activation domains having the requisite functional activities are within the scope of the invention. Additionally, the transactivator fusion protein can include a third polypeptide which promotes transport of the fusion protein to a cell nucleus. For example, a nuclear localization signal (e.g., having an amino acid sequence shown in SEQ ID NO: 5) can be incorporated into the fusion protein. The invention further provides recombinant vectors and host cells comprising nucleic acid encoding a transactivator fusion protein of the invention. The invention still further provides transgenic and homologous recombinant organisms comprising nucleic acid encoding a transactivator fusion protein of the invention.

A transactivator of the invention is used to regulate the transcription of a target transcription unit composed of a nucleotide sequence to be transcribed operatively linked to a minimal promoter sequence and at least one *tet* operator sequence. The nucleotide sequence to be transcribed may encode a protein of interest or an active RNA molecule (e.g., antisense RNA molecule or ribozyme) and may be an exogenous or endogenous nucleotide sequence. A first nucleic acid encoding a transactivator of the invention and a second nucleic acid comprising a target transcription unit for the transactivator can be incorporated into a nucleic acid composition. The nucleic acid composition can be introduced into host cells or organisms (e.g., transgenic and homologous recombinant animals and plants) to allow for tetracycline-inducible expression of the target nucleotide sequence to be transcribed.

In addition to providing a regulatory system for expression of a single nucleotide sequence to be transcribed, the invention also features novel target transcription units that enable coordinate or independent regulation of two or more nucleotide sequences to be transcribed. In one embodiment, coordinate regulation of two nucleotide sequences to be transcribed is achieved using a transcription unit in which a first nucleotide sequence to be transcribed is operatively linked to the 5' end of a *tet* operator sequence(s) and a second nucleotide sequence to be transcribed is operatively linked to the 3' end of the same *tet* operator sequence(s) such that the first and second nucleotide sequences are transcribed in a divergent manner. Suitable bidirectional promoters for such bidirectional transcription units are shown in SEQ ID NOs: 6 and 7. Such transcription units are particularly useful for producing stoichiometric amounts of two subunits of a heterodimeric protein (e.g., antibody chains) in the same cell or for coexpressing a gene of interest and a gene encoding a detectable marker in the same cell, thereby allowing for selection of cells expressing the gene of interest.

In another embodiment, independent regulation of two nucleotide sequences is achieved using a transcription unit in which a first nucleotide sequence to be transcribed is operatively linked to a *tet* operator of a first class type and a second nucleotide sequence to be transcribed is operatively linked to a *tet* operator of a second, different class type. Two different transcriptional activator fusion proteins are then used to regulate the transcription of the two nucleotide sequences: one fusion protein binds to the first class of *tet* operator sequences in the presence of Tc, whereas the other fusion protein binds to the second class of *tet* operator sequences in the absence of Tc (or alternatively, the first fusion protein binds to the first class of *tet* operator sequences in the absence of Tc and the second fusion protein binds to the second class of *tet* operator sequences in the presence of Tc). This transcription unit is particularly useful for independently regulating expression of a therapeutic gene and a suicide gene in a host. For example, expression of the therapeutic gene is stimulated in the presence of tetracycline and then when the therapy is completed, expression of the suicide gene is stimulated by removing the tetracycline.

Another aspect of the invention pertains to methods for stimulating transcription of a nucleotide sequence operatively linked to at least one *tet* operator sequence in a host cell or animal which expresses a transactivator fusion protein of the invention. In a host cell, transcription is stimulated by contacting the cell with Tc or a Tc analogue. In a subject (e.g., transgenic or homologous recombinant organisms), transcription is stimulated by administering Tc or a Tc analogue to the subject. Different Tc analogues, and variation of the concentration of the inducing agent, can be used to modulate the level of induction of gene expression. Preferred Tc analogues for high level gene expression include anhydrotetracycline and doxycycline. The invention further provides a process for producing and isolating proteins of interest using the regulatory system of the invention is also provided.

Another aspect of the invention pertains to tetracycline-regulated transcriptional inhibitors which are useful for inhibiting expression, in a highly controlled manner, of a gene

linked to one or more *tet* operator sequences. The transcriptional inhibitors of the invention comprise a fusion protein composed of at least two polypeptides, a first polypeptide that binds to *tet* operator sequences and a heterologous second polypeptide that directly or indirectly inhibits transcription in eukaryotic cells. The heterologous the second polypeptide is derived from a different protein than the first polypeptide. Because the fusion proteins of the invention include a eukaryotic transcriptional silencer domain, they are anticipated to be more efficient at repressing transcription in eukaryotic cells than is a Tet Repressor alone.

In one embodiment of the invention, the first polypeptide of the inhibitor fusion protein binds to *tet* operator sequences in the absence but not the presence of tetracycline (Tc) or an analogue thereof (e.g., the first polypeptide is preferably a Tet repressor, such as a Tn10-derived Tet repressor having an amino acid sequence shown in SEQ ID NO: 17). In the absence of tetracycline (or Tc analogue), this fusion protein binds to *tet* operator sequences operatively linked to a gene of interest, thereby inhibiting transcription of the gene of interest. In another embodiment, the first polypeptide binds to *tet* operator sequences in the presence but not the absence of tetracycline (e.g., the first polypeptide is preferably a mutated Tet repressor, such as a Tn10-derived Tet repressor having an amino acid substitution at position 71, 95, 101 and/or 102). Preferably, the first polypeptide has an amino acid sequence shown in SEQ ID NO: 19. In the presence of tetracycline (or Tc analogue), this fusion protein binds to *tet* operator sequences operatively linked to a gene of interest, thereby inhibiting transcription of the gene of interest.

The second polypeptide can be a transcriptional "silencer" domain from a protein such as the v-erbA oncogene product, the Drosophila Krueppel protein, the retinoic acid receptor alpha, the thyroid hormone receptor alpha, the yeast Ssn6/Tup1 protein complex, the Drosophila protein even-skipped, SIR1, NeP1, the Drosophila dorsal protein, TSF3, SFI, the Drosophila hunchback protein, the Drosophila knirps protein, WT1, Oct-2.1, the Drosophila engrailed protein, E4BP4 or ZF5. Preferred silencer domains include amino acid residues 403-466 of Krueppel (shown in SEQ ID NO: 21) and amino acid residues 364-635 of v-erbA (shown in SEQ ID NO: 23).

The fusion proteins of the invention may further comprise additional polypeptides, such as a third polypeptide which promotes transport of the fusion protein into a cell nucleus (i.e., a nuclear transport amino acid sequence).

This invention further provides isolated nucleic acid molecules encoding the transcriptional inhibitor fusion proteins of the invention and recombinant expression vectors containing these nucleic acid molecules in a form suitable for expression of the encoded transcriptional inhibitor fusion protein in a host cell. The invention still further provides host cells into which a recombinant expression vectors of the invention has been introduced. Thus, a transcriptional inhibitor fusion protein is expressed in these host cells. The host cell can be, for example, a mammalian cell (e.g., a human cell), a yeast cell, a fungal cell or an insect cell. Moreover, the host cell can be a fertilized non-human oocyte, in which case the

host cell can be used to create a transgenic organism having cells that express the transcriptional inhibitor fusion protein. Still further, the recombinant expression vector can be designed to allow homologous recombination between the nucleic acid encoding the fusion protein and a target gene in a host cell. Such homologous recombination vectors can
5 be used to create homologous recombinant animals that express a fusion protein of the invention.

In a preferred embodiment, the host cells (or cells of a host organism) also contain a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence (e.g., a gene of interest whose expression can be regulated by Tc or an analogue thereof). To
10 regulate transcription of the *tet* operator-linked gene of interest in these host cells, the concentration of Tc (or analogue thereof) in contact with the host cell is altered. For example, when the transcriptional inhibitor fusion protein binds to *tet* operator sequences in the absence of Tc, the concentration of Tc in contact with the cells is decreased to thereby inhibit transcription of the *tet* operator-linked gene of interest (e.g., if cells are first cultured
15 in the presence of Tc, then Tc can be removed from the culture medium to inhibit transcription of the gene of interest). Alternatively, when the transcriptional inhibitor fusion protein binds to *tet* operator sequences in the presence of Tc, the concentration of Tc in contact with the cells is increased to thereby inhibit transcription of the *tet* operator-linked gene of interest (e.g., if cells are first cultured in the absence of Tc, then Tc can be added to
20 the culture medium to inhibit transcription of the gene of interest).

The transcriptional inhibitor fusion proteins of the invention are useful for inhibiting gene expression in a variety of situations, as described further herein. In a particularly preferred embodiment, transcription of a *tet* operator(*tetO*)-linked gene of interest is regulated by a combination of tetracycline-regulated transcriptional inhibitor and activator fusion
25 proteins in the same host cell to allow for precise control of the expression level of the gene of interest. For example, an activator fusion protein that binds to *tetO* only in the presence of Tc and an inhibitor fusion protein that binds to *tetO* only in the absence of Tc are expressed in a host cell that contains a *tetO*-linked gene of interest. In the absence of Tc, basal levels of transcription of the gene of interest are inhibited by the inhibitor fusion protein. Upon
30 contact of the cell with Tc (or analogue), transcription of the gene of interest is stimulated by the activator fusion protein. The activator and inhibitor fusion proteins of the invention can also be used in combination to regulate the expression of multiple *tetO*-linked genes of interest.

Novel kits for regulating the expression of a gene of interest are also within the scope
35 of the invention. In one embodiment, the kits of the invention can include at least one nucleic acid molecule encoding a transactivator fusion protein of the invention and a target transcription unit into which a gene of interest can be cloned such that the gene is operatively linked to a *tet* operator sequence(s). Alternatively or additionally, the kits can include a nucleic acid molecule encoding a transcriptional inhibitor fusion protein of the invention.

Target transcription units into which multiple genes of interest can be cloned, e.g., for coordinate or independent regulation, can also be included in kits of the invention. Moreover, at least one tetracycline or tetracycline analogue may be included in the kits.

5 **Brief Description of the Drawings**

Figure 1 is a bar graph depicting the stimulation of luciferase activity in HR5-C11 cells by tetracycline and different tetracycline analogues (1 µg/ml f.c.). Cells were grown in the absence (-) or presence of the indicated tetracyclines for 3 days before luciferase activity was determined. Each solid and hatched bar represents the luciferase activity of a single
10 culture dish.

Figure 2 is a graph depicting the relative luciferase activity in HR5-C11 cells when incubated with different concentrations of doxycycline. The results of three independent experiments are shown.

Figure 3 is a graph depicting the kinetics of induction of luciferase activity in HR5-
15 C11 cells by doxycycline. HR5-C11 cultures were exposed to 1 µg/ml of doxycycline and luciferase activity was measured after different time intervals; (•) cultures containing doxycycline, (o) cultures grown in the absence of antibiotic.

Figure 4 shows the amino acid sequences of various classes of Tet repressors, illustrating the homology between the amino acid sequences of different classes of Tet
20 repressors, as compared to class B Tet repressors (e.g., Tn10-derived). Amino acid positions in other classes of Tet repressors that are identical to class B are indicated by a dash.

Figure 5 shows the nucleotide sequences of *tet* operators of different classes: class A (SEQ ID NO: 11), class B (SEQ ID NO: 12), class C (SEQ ID NO: 13), class D (SEQ ID NO: 14) and class E (SEQ ID NO: 15).

Figure 6 is a schematic diagram of a bidirectional promoter construct for coordinate
25 regulation of two genes of interest operatively linked to the same *tet* operators for regulation by a tetracycline-regulated transcriptional activator.

Figure 7A (SEQ ID NO: 6) shows the nucleotide sequence of a bidirectional promoter
30 region for coordinate regulation of two genes of interest by a tetracycline-regulated transcriptional activator.

Figure 7B (SEQ ID NO: 7) shows the nucleotide sequence of a bidirectional promoter
region for coordinate regulation of two genes of interest by a tetracycline-regulated transcriptional activator.

Figure 8 is two graphs depicting coordinate expression of luciferase and β-
35 galactosidase activity by a tetracycline-regulated transcriptional activator.

Figure 9A-B are schematic diagrams of self-regulating promoters for expression of tetracycline-regulated transcriptional activators (tTA). Panel A illustrates self-regulation of expression of a wild-type Tet repressor-containing transactivator fusion protein that binds to *tet* operators in the absence of Tc. Panel B illustrates self-regulation of expression of a

mutated Tet repressor-containing transactivator fusion protein that binds to *tet* operators in the presence of Tc.

5 *Figure 10* is a schematic diagram of the negative and positive regulation of a *tet* operator (*tetOwt*)-linked gene of interest by a tetracycline-regulated transcriptional inhibitor protein (tSD) and a tetracycline-inducible transcriptional activator fusion protein (rtTA), respectively, in the presence of increasing concentrations of the tetracycline analogue doxycycline.

10 *Figure 11* is a schematic diagram of the construction of TetR-silencer domain fusion constructs by in-frame fusion of nucleic acid encoding either a Krueppel or v-erbA silencer domain to the 3' end of nucleic acid encoding a Tet repressor (*tetR* gene).

15 *Figure 12* is a graphic representation of the expression of luciferase activity in mice transgenic for the luciferase reporter gene alone (checked columns at right) or double transgenic animals carrying the luciferase reporter gene and a tTA^R transgene, either in the absence of doxycycline (dark columns in middle) or in the presence of doxycycline (light columns at left).

Detailed Description of the Invention

20 This invention pertains to nucleic acid molecules and proteins which can be used to regulate the expression of genes in eukaryotic cells or animals in a highly controlled manner. Regulation of gene expression by the system of the invention involves at least two components: A gene which is operatively linked to a regulatory sequence and a protein which, in either the presence or absence of an inducible agent, binds to the regulatory sequence and either activates or inhibits transcription of the gene. The system of the invention utilizes components of the Tet repressor/operator/inducer system of prokaryotes to
25 stimulate gene expression in eukaryotic cells.

Various aspects of the invention pertain to fusion proteins which are capable of either activating or inhibiting gene transcription when bound to *tet* operator (*tetO*) sequences, but which bind to *tet* operator sequences only in the presence or, alternatively, in the absence of tetracycline, or an analogue thereof. Thus, in a host cell, transcription of a gene operatively
30 linked to a *tet* operator sequence(s) is stimulated or inhibited by a fusion protein of the invention by altering the concentration of tetracycline (or analogue) in contact with the host cell (e.g., adding or removing tetracycline from a culture medium, or administering or ceasing to administer tetracycline to a host organism, etc.).

35 The invention further pertains to target transcription units for regulation by the fusion protein of the invention. In addition to allowing for regulation of a single *tet*-operator linked gene of interest, the invention also provides novel transcription units containing two or more genes to be transcribed that can be regulated in either a coordinate or independent manner by a transactivator fusion protein of the invention. Methods for stimulating or inhibiting transcription of a gene using tetracycline (or analogues thereof), and kits which contain the

components of the regulatory system described herein, are also encompassed by the invention.

In the following subsections, the nucleic acids and proteins comprising the components of the inducible regulatory system of the invention, and their interrelationship, are discussed in greater detail. The subsections are as follows:

I. Tetracycline-Inducible Transcriptional Activators

- A. The first polypeptide of the transactivator fusion protein
- B. The second polypeptide of the transactivator fusion protein
- 10 C. A third polypeptide of the transactivator fusion protein

II. Expression of a Transactivator Fusion Protein

- A. Expression vectors
- B. Host cells
- 15 C. Introduction of nucleic acid into host cells
- D. Transgenic Organisms
- E. Homologous Recombinant Organisms

III. Target Transcription Units Regulated by a Tetracycline-Inducible Transactivator

- 20 A. Regulation of expression of tet operator-linked nucleotide sequences
- B. Coordinate regulation of two nucleotide sequences
- C. Independent regulation of multiple nucleotide sequences
- D. Combined coordinate and independent regulation of multiple nucleotide sequences

25

IV. Tetracycline-Regulated Transcriptional Inhibitors

- A. The first polypeptide of the transcriptional inhibitor fusion protein
- B. The second polypeptide of the transcriptional inhibitor fusion protein
- C. A third polypeptide of the transcriptional inhibitor fusion protein
- 30 D. Expression of the transcriptional inhibitor fusion protein

V. Kits of the Invention

VI. Regulation of Gene Expression by Tetracycline or Analogues Thereof

- 35 A. Stimulation of gene expression by transactivator fusion proteins
- B. Inhibition of gene expression by transcriptional inhibitor fusion proteins
- C. Combined positive and negative regulation of gene expression

VII. Applications of the Invention

- A. Gene Therapy
- B. Production of Proteins in Vitro
- C. Production of Proteins in Vivo
- 5 D. Animal Models of Human Disease
- E. Production of Stable Cell Lines for Cloning

I. Tetracycline-Inducible Transcriptional Activators

In the inducible regulatory system of the invention, transcription of a gene is activated
10 by a transcriptional activator protein, also referred to herein simply as a transactivator. The transactivator of the invention is a fusion protein. One aspect of the invention thus pertains to fusion proteins and nucleic acids (e.g., DNA) encoding fusion proteins. The term "fusion protein" is intended to describe at least two polypeptides, typically from different sources, which are operatively linked. With regard to the polypeptides, the term "operatively linked"
15 is intended to mean that the two polypeptides are connected in manner such that each polypeptide can serve its intended function. Typically, the two polypeptides are covalently attached through peptide bonds. The fusion protein is preferably produced by standard recombinant DNA techniques. For example, a DNA molecule encoding the first polypeptide is ligated to another DNA molecule encoding the second polypeptide, and the resultant hybrid
20 DNA molecule is expressed in a host cell to produce the fusion protein. The DNA molecules are ligated to each other in a 5' to 3' orientation such that, after ligation, the translational frame of the encoded polypeptides is not altered (i.e., the DNA molecules are ligated to each other in-frame).

25 *A. The first polypeptide of the transactivator fusion protein*

The transactivator fusion protein of the invention is composed, in part, of a first polypeptide which binds to a *tet* operator sequence in the presence of tetracycline (Tc), or an analogue thereof. The first polypeptide of the fusion protein is preferably a mutated Tet repressor. The term "mutated Tet repressor" is intended to include polypeptides having an
30 amino acid sequence which is similar to a wild-type Tet repressor but which has at least one amino acid difference from the wild-type Tet repressor. The term "wild-type Tet repressor" is intended to describe a protein occurring in nature which represses transcription from *tet* operator sequences in prokaryotic cells in the absence of Tc. The amino acid difference(s) between a mutated Tet repressor and a wild-type Tet repressor may be substitution of one or
35 more amino acids, deletion of one or more amino acids or addition of one or more amino acids. The mutated Tet repressor of the invention has the following functional properties: 1) the polypeptide can bind to a *tet* operator sequence, i.e., it retains the DNA binding specificity of a wild-type Tet repressor; and 2) it is regulated in a reverse manner by

tetracycline than a wild-type Tet repressor, i.e., the mutated Tet repressor binds to a *ter* operator sequence only the presence of Tc (or Tc analogue) rather than in the absence of Tc.

In a preferred embodiment, a mutated Tet repressor having the functional properties described above is created by substitution of amino acid residues in the sequence of a wild-type Tet repressor. For example, as described in Example 1, a Tn10-derived Tet repressor having amino acid substitutions at amino acid positions 71, 95, 101 and 102 has the desired functional properties and thus can be used as the first polypeptide in the transactivator fusion protein of the invention. The amino acid sequence of this mutated Tet repressor is shown in SEQ ID NO: 2 (positions 1-207). In one embodiment of the mutated Tet repressor, position 71 is mutated from glutamic acid to lysine, position 95 is mutated from aspartic acid to asparagine, position 101 is mutated from leucine to serine and position 102 is mutated from glycine to aspartic acid, although the invention is not limited to these particular mutations. Mutation of fewer than all four of these amino acid positions may be sufficient to achieve a Tet repressor with the desired functional properties. Accordingly, a Tet repressor is preferably mutated at at least one of these positions. Other amino acid substitutions, deletions or additions at these or other amino acid positions which retain the desired functional properties of the mutated Tet repressor are within the scope of the invention. The crystal structure of a Tet repressor-tetracycline complex, as described in Hinrichs, W. et al. (1994) *Science* 264:418-420, can be used for rational design of mutated Tet repressors. Based upon this structure, amino acid position 71 is located outside the tetracycline binding pocket, suggesting mutation at this site may not be necessary to achieve the desired functional properties of a mutated Tet repressor of the invention. In contrast, amino acid positions 95, 101 and 102 are located within the conserved tetracycline binding pocket. Thus, the tetracycline binding pocket of a Tet repressor may be targeted for mutation to create a mutated Tet repressor of the invention.

Additional mutated Tet repressors for incorporation into a fusion protein of the invention can be created according to the teachings of the invention. A number of different classes of Tet repressors have been described, e.g., A, B, C, D and E (of which the Tn10-encoded repressor is a class B repressor). The amino acid sequences of the different classes of Tet repressors share a high degree of homology (i.e., 40-60 % across the length of the proteins), including in the region encompassing the above-described mutations. The amino acid sequences of various classes of Tet repressors are shown and compared in Figure 4, and are also described in Tovar, K. et al. (1988) *Mol. Gen. Genet.* 215:76-80. Accordingly, equivalent mutations to those described above for the Tn10-derived Tet repressor can be made in other classes of Tet repressors for inclusion in a fusion protein of the invention. For example, amino acid position 95, which is an aspartic acid in all five repressor classes, can be mutated to asparagine in any class of repressor. Similarly, position 102, which is glycine in all five repressor classes, can be mutated to aspartic acid in any class of repressor.

Additional suitable equivalent mutations will be apparent to those skilled in the art and can be

created and tested for functionality by procedures described herein. Nucleotide and amino acid sequences of Tet repressors of the A, C, D and E classes are disclosed in Waters, S.H. et al. (1983) *Nucl. Acids Res* 11:6089-6105, Unger, B. et al. (1984) *Gene* 31: 103-108, Unger, B. et al. (1984) *Nucl Acids Res.* 12:7693-7703 and Tovar, K. et al. (1988) *Mol. Gen. Genet.* 215:76-80, respectively. These wild-type sequences can be mutated according to the teachings of the invention for use in the inducible regulatory system described herein.

Alternative to the above-described mutations, additional suitable mutated Tet repressors (i.e., having the desired functional properties described above) can be created by mutagenesis of a wild type Tet repressor and selection as described in Example 1. The nucleotide and amino acid sequences of wild-type class B Tet repressors are disclosed in Hillen, W. and Schollmeier, K. (1983) *Nucl. Acids Res.* 11:525-539 and Postle, K. et al. (1984) *Nucl. Acids Res.* 12:4849-4863. The nucleotide and amino acid sequences of wild-type class A, C, D and E type repressors are cited above. A mutated Tet repressor can be created and selected, for example as follows: a nucleic acid (e.g., DNA) encoding a wild-type Tet repressor is subjected to random mutagenesis and the resultant mutated nucleic acids are incorporated into an expression vector and introduced into a host cell for screening. A screening assay is used which allows for selection of a Tet repressor which binds to a *tet* operator sequence only in the presence of tetracycline. For example, a library of mutated nucleic acids in an expression vector can be introduced into an *E. coli* strain in which *tet* operator sequences control the expression of a gene encoding a Lac repressor and the Lac repressor controls the expression of a gene encoding an selectable marker (e.g., drug resistance). Binding of a Tet repressor to *tet* operator sequences in the bacteria will inhibit expression of the Lac repressor, thereby inducing expression of the selectable marker gene. Cells expressing the marker gene are selected based upon the selectable phenotype (e.g., drug resistance). For wild-type Tet repressors, expression of the selectable marker gene will occur in the absence of Tc. A nucleic acid encoding a mutated Tet repressor is selected using this system based upon the ability of the nucleic acid to induce expression of the selectable marker gene in the bacteria only in the presence of Tc.

A first polypeptide of the transactivator fusion protein (e.g., the mutated Tet repressor) has the property of binding specifically to a *tet* operator sequence. Each class of Tet repressor has a corresponding target *tet* operator sequence. Accordingly, the term "*tet* operator sequence" is intended to encompass all classes of *tet* operator sequences, e.g. class A, B, C, D, and E. Nucleotide sequences of these five classes of *tet* operators are shown in Figure 5 and SEQ ID NOs: 11-15, and are described in Waters, S.H. et al. (1983) cited *supra*, Hillen, W. and Schollenmeier, K. (1983) cited *supra*, Stüber, D. and Bujard, H. (1981) *Proc. Natl. Acad. Sci. USA* 78:167-171, Unger, B. et al. (1984) cited *supra* and Tovar, K. et al. (1988) cited *supra*. In a preferred embodiment, the mutated Tet repressor is a Tn10-encoded repressor (i.e., class B) and the *tet* operator sequence is a class B *tet* operator sequence.

Alternatively, a mutated class A Tet repressor can be used with a class A *tet* operator sequence, and so on for the other classes of Tet repressor/operators.

Another approach for creating a mutated Tet repressor which binds to a class A *tet* operator is to further mutate the already mutated Tn10-derived Tet repressor described herein (a class B repressor) such that it no longer binds efficiently to a class B type operator but instead binds efficiently to a class A type operator. It has been found that nucleotide position 6 of class A or B type operators is the critical nucleotide for recognition of the operator by its complementary repressor (position 6 is a G/C pair in class B operators and an A/T pair in class A operators) (see Wissman et al. (1988) *J. Mol. Biol.* 202:397-406). It has also been found that amino acid position 40 of a class A or class B Tet repressor is the critical amino acid residue for recognition of position 6 of the operator (amino acid position 40 is a threonine in class B repressors but is an alanine in class A repressors). It still further has been found that substitution of Thr40 of a class B repressor with Ala alters its binding specificity such that the repressor can now bind a class A operator (similarly, substitution of Ala40 of a class A repressor with Thr alters its binding specificity such that the repressor can now bind a class B operator) (see Altschmied et al. (1988) *EMBO J.* 7:4011-4017). Accordingly, one can alter the binding specificity of the mutated Tn10-derived Tet repressor disclosed herein by additionally changing amino acid residue 40 from Thr to Ala by standard molecular biology techniques (e.g., site directed mutagenesis).

A mutated Tet repressor having specific mutations (e.g., at positions 71, 95, 101 and/or 102, as described above) can be created by introducing nucleotide changes into a nucleic acid encoding a wild-type repressor by standard molecular biology techniques, e.g. site directed mutagenesis or PCR-mediated mutagenesis using oligonucleotide primers incorporating the nucleotide mutations. Alternatively, when a mutated Tet repressor is identified by selection from a library, the mutated nucleic acid can be recovered from the library vector. To create a transactivator fusion protein of the invention, a nucleic acid encoding a mutated Tet repressor is then ligated in-frame to another nucleic acid encoding a transcriptional activation domain and the fusion construct is incorporated into a recombinant expression vector. The transactivator fusion protein can be expressed by introducing the recombinant expression vector into a host cell or animal.

B. The second polypeptide of the transactivator fusion protein

The first polypeptide of the transactivator fusion protein is operatively linked to a second polypeptide which directly or indirectly activates transcription in eukaryotic cells. To operatively link the first and second polypeptides, typically nucleotide sequences encoding the first and second polypeptides are ligated to each other in-frame to create a chimeric gene encoding a fusion protein, although the first and second polypeptides can be operatively linked by other means that preserve the function of each polypeptide (e.g., chemically crosslinked). In a preferred embodiment, the second polypeptide of the transactivator itself

possesses transcriptional activation activity (i.e., the second polypeptide directly activates transcription). In another embodiment, the second polypeptide activates transcription by an indirect mechanism, through recruitment of a transcriptional activation protein to interact with the fusion protein. Accordingly, the term "a polypeptide which activates transcription in eukaryotic cells" as used herein is intended to include polypeptides which either directly or indirectly activates transcription.

Polypeptides which can function to activate transcription in eukaryotic cells are well known in the art. In particular, transcriptional activation domains of many DNA binding proteins have been described and have been shown to retain their activation function when the domain is transferred to a heterologous protein. A preferred polypeptide for use in the fusion protein of the invention is the herpes simplex virus virion protein 16 (referred to herein as VP16, the amino acid sequence of which is disclosed in Triezenberg, S.J. et al. (1988) *Genes Dev.* 2:718-729). In one embodiment, about 127 of the C-terminal amino acids of VP16 are used. For example, a polypeptide having an amino acid sequence shown in SEQ ID NO: 2 (positions 208-335) can be used as the second polypeptide in the fusion protein. In another embodiment, at least one copy of about 11 amino acids from the C-terminal region of VP16 which retain transcriptional activation ability is used as the second polypeptide. Preferably, a dimer of this region (i.e., about 22 amino acids) is used. Suitable C-terminal peptide portions of VP16 are described in Seipel, K. et al. (*EMBO J.* (1992) 13:4961-4968). For example, a dimer of a peptide having an amino acid sequence shown in SEQ ID NO: 4 (encoded by a nucleotide sequence shown in SEQ ID NO: 3) can be used as the second polypeptide in the fusion protein.

Other polypeptides with transcriptional activation ability in eukaryotic cells can be used in the fusion protein of the invention. Transcriptional activation domains found within various proteins have been grouped into categories based upon similar structural features. Types of transcriptional activation domains include acidic transcription activation domains, proline-rich transcription activation domains, serine/threonine-rich transcription activation domains and glutamine-rich transcription activation domains. Examples of acidic transcriptional activation domains include the VP16 regions already described and amino acid residues 753-881 of GAL4. Examples of proline-rich activation domains include amino acid residues 399-499 of CTF/NF1 and amino acid residues 31-76 of AP2. Examples of serine/threonine-rich transcription activation domains include amino acid residues 1-427 of ITF1 and amino acid residues 2-451 of ITF2. Examples of glutamine-rich activation domains include amino acid residues 175-269 of Oct1 and amino acid residues 132-243 of Sp1. The amino acid sequences of each of the above described regions, and of other useful transcriptional activation domains, are disclosed in Seipel, K. et al. (*EMBO J.* (1992) 13:4961-4968).

In addition to previously described transcriptional activation domains, novel transcriptional activation domains, which can be identified by standard techniques, are within

the scope of the invention. The transcriptional activation ability of a polypeptide can be assayed by linking the polypeptide to another polypeptide having DNA binding activity and determining the amount of transcription of a target sequence that is stimulated by the fusion protein. For example, a standard assay used in the art utilizes a fusion protein of a putative
5 transcriptional activation domain and a GAL4 DNA binding domain (e.g., amino acid residues 1-93). This fusion protein is then used to stimulate expression of a reporter gene linked to GAL4 binding sites (see e.g., Seipel, K. et al. (1992) *EMBO J.* 11:4961-4968 and references cited therein).

In another embodiment, the second polypeptide of the fusion protein indirectly
10 activates transcription by recruiting a transcriptional activator to interact with the fusion protein. For example, a mutated tetR of the invention can be fused to a polypeptide domain (e.g., a dimerization domain) capable of mediating a protein-protein interaction with a transcriptional activator protein, such as an endogenous activator present in a host cell. It has been demonstrated that functional associations between DNA binding domains and
15 transactivation domains need not be covalent (see e.g., Fields and Song (1989) *Nature* 340:245-247; Chien et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:9578-9582; Gyuris et al. (1993) *Cell* 75:791-803; and Zervos, A.S. (1993) *Cell* 72:223-232). Accordingly, the second polypeptide of the fusion protein may not directly activate transcription but rather may form a stable interaction with an endogenous polypeptide bearing a compatible protein-protein
20 interaction domain and transactivation domain. Examples of suitable interaction (or dimerization) domains include leucine zippers (Landschulz et al. (1989) *Science* 243:1681-1688), helix-loop-helix domains (Murre, C. et al. (1989) *Cell* 58:537-544) and zinc finger domains (Frankel, A.D. et al. (1988) *Science* 240:70-73). Interaction of a dimerization domain present in the fusion protein with an endogenous nuclear factor results in
25 recruitment of the transactivation domain of the nuclear factor to the fusion protein, and thereby to a *tet* operator sequence to which the fusion protein is bound.

C. A third polypeptide of the transactivator fusion protein

In addition to a mutated Tet repressor and a transcriptional activation domain, a fusion
30 protein of the invention can contain an operatively linked third polypeptide which promotes transport of the fusion protein to a cell nucleus. Amino acid sequences which, when included in a protein, function to promote transport of the protein to the nucleus are known in the art and are termed nuclear localization signals (NLS). Nuclear localization signals typically are composed of a stretch of basic amino acids. When attached to a heterologous protein (e.g., a
35 fusion protein of the invention), the nuclear localization signal promotes transport of the protein to a cell nucleus. The nuclear localization signal is attached to a heterologous protein such that it is exposed on the protein surface and does not interfere with the function of the protein. Preferably, the NLS is attached to one end of the protein, e.g. the N-terminus. The amino acid sequence of a non-limiting example of an NLS that can be included in a fusion

protein of the invention is shown in SEQ ID NO: 5. Preferably, a nucleic acid encoding the nuclear localization signal is spliced by standard recombinant DNA techniques in-frame to the nucleic acid encoding the fusion protein (e.g., at the 5' end).

5 The plasmid pUHD17-1 (described in further detail in Example 1), which comprises a transactivator of the invention having the nucleotide sequence shown in SEQ ID NO: 1, has been deposited on July 8, 1994 under the provisions of the Budapest Treaty at the Deutsche Sammlung Von Mikroorganismen und ZellKulturen GmbH (DSMZ) in Braunschweig, Germany and assigned deposit number DSM 9279.

10

II. Expression of a Transactivator Fusion Protein

A. Expression Vectors

15 A nucleic acid of the invention encoding a transactivator fusion protein, as described above, can be incorporated into a recombinant expression vector in a form suitable for expression of the fusion protein in a host cell. The term "in a form suitable for expression of the fusion protein in a host cell" is intended to mean that the recombinant expression vector includes one or more regulatory sequences operatively linked to the nucleic acid encoding the fusion protein in a manner which allows for transcription of the nucleic acid into mRNA and
20 translation of the mRNA into the fusion protein. The term "regulatory sequence" is art-recognized and intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are known to those skilled in the art and are described in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). It should be understood that the
25 design of the expression vector may depend on such factors as the choice of the host cell to be transfected and/or the amount of fusion protein to be expressed.

 When used in mammalian cells, a recombinant expression vector's control functions are often provided by viral genetic material. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. Use of viral
30 regulatory elements to direct expression of the fusion protein can allow for high level constitutive expression of the fusion protein in a variety of host cells. In a preferred recombinant expression vector, the sequences encoding the fusion protein are flanked upstream (i.e., 5') by the human cytomegalovirus IE promoter and downstream (i.e., 3') by an SV40 poly(A) signal. For example, an expression vector similar to that described in Example
35 1 can be used. The human cytomegalovirus IE promoter is described in Boshart et al. (1985) *Cell* 41:521-530. Other ubiquitously expressing promoters which can be used include the HSV-Tk promoter (disclosed in McKnight et al. (1984) *Cell* 37:253-262) and β -actin promoters (e.g., the human β -actin promoter as described by Ng et al. (1985) *Mol. Cell. Biol.* 5:2720-2732).

Alternatively, the regulatory sequences of the recombinant expression vector can direct expression of the fusion protein preferentially in a particular cell type, i.e., tissue-specific regulatory elements can be used. Non-limiting examples of tissue-specific promoters which can be used include the albumin promoter (liver-specific; Pinkert et al. (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji et al. (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund et al. (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

Alternatively, a self-regulating construct encoding a transactivator fusion protein can be created. To accomplish this, nucleic acid encoding the fusion protein is operatively linked to a minimal promoter sequence and at least one *tet* operator sequence. For example, the nucleic acid of SEQ ID NO: 1 can be linked to a promoter having a nucleotide sequence shown in SEQ ID NO: 8, 9 or 10 (the nucleic acids of SEQ ID NOs: 8 and 9 comprise a minimal CMV promoter and ten *tet* operators; the nucleic acids of SEQ ID NO: 10 comprises a TK promoter and ten *tet* operators). A schematic diagram of such a self-regulating construct is shown in Figure 9B. When this nucleic acid is introduced into a cell (e.g., in a recombinant expression vector), a small amount of basal transcription of the transactivator gene is likely to occur due to "leakiness". In the presence of Tc (or analogue thereof) this small amount of the transactivator fusion protein will bind to the *tet* operator sequence(s) upstream of the nucleotide sequence encoding the transactivator and stimulate additional transcription of the nucleotide sequence encoding the transactivator, thereby leading to further production of the transactivator fusion protein in the cell. It will be appreciated by those skilled in the art that such a self-regulating promoter can also be used in conjunction with other tetracycline-regulated transactivators, such as the wild-type Tet repressor fusion protein (tTA) described in Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551, which binds to *tet* operators in the absence of Tc (as illustrated in Figure 9A). When used in conjunction with this transactivator, self-regulated transcription of the nucleotide sequence encoding this transactivator is stimulated in the absence of Tc. The plasmid pUHD15-3, which comprises nucleotide sequences encoding the tTA described in Gossen and Bujard (1992), cited *supra*, operatively linked to a self-regulating promoter, has been deposited on July 8, 1994 under the provisions of the Budapest Treaty at the Deutsche Sammlung Von Mikroorganismen und ZellKulturen GmbH (DSM) in Braunschweig, Germany and assigned deposit number DSM 9280.

In one embodiment, the recombinant expression vector of the invention is a plasmid, such as that described in Example 1. Alternatively, a recombinant expression vector of the invention can be a virus, or portion thereof, which allows for expression of a nucleic acid introduced into the viral nucleic acid. For example, replication defective retroviruses, adenoviruses and adeno-associated viruses can be used. Protocols for producing recombinant retroviruses and for infecting cells *in vitro* or *in vivo* with such viruses can be found in Current Protocols in Molecular Biology, Ausubel, F.M. et al. (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are well known to those skilled in the art. Examples of suitable packaging virus lines include ψ Crip, ψ Cre, ψ 2 and ψ Am. The genome of adenovirus can be manipulated such that it encodes and expresses a transactivator fusion protein but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. See for example Berkner et al. (1988) *BioTechniques* 6:616; Rosenfeld et al. (1991) *Science* 252:431-434; and Rosenfeld et al. (1992) *Cell* 68:143-155. Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or other strains of adenovirus (e.g., Ad2, Ad3, Ad7 etc.) are well known to those skilled in the art. Alternatively, an adeno-associated virus vector such as that described in Tratschin et al. (1985) *Mol. Cell. Biol.* 5:3251-3260 can be used to express a transactivator fusion protein.

20 B. Host Cells

A fusion protein of the invention is expressed in a eukaryotic cell by introducing nucleic acid encoding the fusion protein into a host cell, wherein the nucleic acid is in a form suitable for expression of the fusion protein in the host cell. For example, a recombinant expression vector of the invention, encoding the fusion protein, is introduced into a host cell. Alternatively, nucleic acid encoding the fusion protein which is operatively linked to regulatory sequences (e.g., promoter sequences) but without additional vector sequences can be introduced into a host cell. As used herein, the term "host cell" is intended to include any eukaryotic cell or cell line so long as the cell or cell line is not incompatible with the protein to be expressed, the selection system chosen or the fermentation system employed. Non-limiting examples of mammalian cell lines which can be used include CHO dhfr⁻ cells (Urlaub and Chasin (1980) *Proc. Natl. Acad. Sci. USA* 77:4216-4220), 293 cells (Graham et al. (1977) *J. Gen. Virol.* 36: pp59) or myeloma cells like SP2 or NS0 (Galfre and Milstein (1981) *Meth. Enzymol.* 73(B):3-46).

In addition to cell lines, the invention is applicable to normal cells, such as cells to be modified for gene therapy purposes or embryonic cells modified to create a transgenic or homologous recombinant animal. Examples of cell types of particular interest for gene therapy purposes include hematopoietic stem cells, myoblasts, hepatocytes, lymphocytes, neuronal cells and skin epithelium and airway epithelium. Additionally, for transgenic or homologous recombinant animals, embryonic stem cells and fertilized oocytes can be

modified to contain nucleic acid encoding a transactivator fusion protein. Moreover, plant cells can be modified to create transgenic plants.

The invention is broadly applicable and encompasses non-mammalian eukaryotic cells as well, including insect (e.g., *Sp. frugiperda*), yeast (e.g., *S. cerevisiae*, *S. pombe*, *P. pastoris*, *K. lactis*, *H. polymorpha*; as generally reviewed by Fleer, R. (1992) *Current Opinion in Biotechnology* 3(5):486-496), fungal and plant cells. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA). The fusion protein can be expressed in insect cells using baculovirus expression vectors (e.g., as described in O'Reilly *et al.* (1992) *Baculovirus Expression Vectors: A Laboratory Manual*, Stockton Press). Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., SF 9 cells) include the pAc series (Smith *et al.*, (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow, V.A., and Summers, M.D., (1989) *Virology* 170:31-39).

15

C. Introduction of Nucleic Acid into a Host Cell

Nucleic acid encoding the fusion protein can be introduced into a host cell by standard techniques for transfecting eukaryotic cells. The term "transfecting" or "transfection" is intended to encompass all conventional techniques for introducing nucleic acid into host cells, including calcium phosphate co-precipitation, DEAE-dextran-mediated transfection, lipofection, electroporation and microinjection. Suitable methods for transfecting host cells can be found in Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks.

The number of host cells transformed with a nucleic acid of the invention will depend, at least in part, upon the type of recombinant expression vector used and the type of transfection technique used. Nucleic acid can be introduced into a host cell transiently, or more typically, for long term regulation of gene expression, the nucleic acid is stably integrated into the genome of the host cell or remains as a stable episome in the host cell. Plasmid vectors introduced into mammalian cells are typically integrated into host cell DNA at only a low frequency. In order to identify these integrants, a gene that contains a selectable marker (e.g., drug resistance) is generally introduced into the host cells along with the nucleic acid of interest. Preferred selectable markers include those which confer resistance to certain drugs, such as G418 and hygromycin. Selectable markers can be introduced on a separate plasmid from the nucleic acid of interest or, are introduced on the same plasmid. Host cells transfected with a nucleic acid of the invention (e.g., a recombinant expression vector) and a gene for a selectable marker can be identified by selecting for cells using the selectable marker. For example, if the selectable marker encodes a gene conferring neomycin resistance, host cells which have taken up nucleic acid can be selected with G418. Cells that have incorporated the selectable marker gene will survive, while the other cells die.

A host cell transfected with a nucleic acid encoding a fusion protein of the invention can be further transfected with one or more nucleic acids which serve as the target for the fusion protein. The target nucleic acid comprises a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence (described in more detail in Section III below).

Nucleic acid encoding the fusion protein of the invention can be introduced into eukaryotic cells growing in culture *in vitro* by conventional transfection techniques (e.g., calcium phosphate precipitation, DEAE-dextran transfection, electroporation etc.). Nucleic acid can also be transferred into cells *in vivo*, for example by application of a delivery mechanism suitable for introduction of nucleic acid into cells *in vivo*, such as retroviral vectors (see e.g., Ferry, N et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8377-8381; and Kay, M.A. et al. (1992) *Human Gene Therapy* 3:641-647), adenoviral vectors (see e.g., Rosenfeld, M.A. (1992) *Cell* 68:143-155; and Herz, J. and Gerard, R.D. (1993) *Proc. Natl. Acad. Sci. USA* 90:2812-2816), receptor-mediated DNA uptake (see e.g., Wu, G. and Wu, C.H. (1988) *J. Biol. Chem.* 263:14621; Wilson et al. (1992) *J. Biol. Chem.* 267:963-967; and U.S. Patent No. 5,166,320), direct injection of DNA (see e.g., Acsadi et al. (1991) *Nature* 332: 815-818; and Wolff et al. (1990) *Science* 247:1465-1468) or particle bombardment (see e.g., Cheng, L. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:4455-4459; and Zelenin, A.V. et al. (1993) *FEBS Letters* 315:29-32). Thus, for gene therapy purposes, cells can be modified *in vitro* and administered to a subject or, alternatively, cells can be directly modified *in vivo*.

D. Transgenic Organisms

Nucleic acid a transactivator fusion protein can transferred into a fertilized oocyte of a non-human animal to create a transgenic animal which expresses the fusion protein of the invention in one or more cell types. A transgenic animal is an animal having cells that contain a transgene, wherein the transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic, stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. In one embodiment, the non-human animal is a mouse, although the invention is not limited thereto. In other embodiments, the transgenic animal is a goat, sheep, pig, cow or other domestic farm animal. Such transgenic animals are useful for large scale production of proteins (so called "gene pharming").

A transgenic animal can be created, for example, by introducing a nucleic acid encoding the fusion protein (typically linked to appropriate regulatory elements, such as a constitutive or tissue-specific enhancer) into the male pronuclei of a fertilized oocyte, e.g., by microinjection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Intronic sequences and polyadenylation signals can also be included in the transgene to

increase the efficiency of expression of the transgene. Methods for generating transgenic animals, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009 and Hogan, B. et al., (1986) *A Laboratory Manual*, Cold Spring Harbor, New York, Cold Spring Harbor

5 Laboratory. A transgenic founder animal can be used to breed additional animals carrying the transgene. Transgenic animals carrying a transgene encoding the fusion protein of the invention can further be bred to other transgenic animals carrying other transgenes, e.g., to a transgenic animal which contains a gene operatively linked to a *tet* operator sequence (discussed in more detail in Section III below).

10 It will be appreciated that, in addition to transgenic animals, the regulatory system described herein can be applied to other transgenic organisms, such as transgenic plants. Transgenic plants can be made by conventional techniques known in the art. Accordingly, the invention encompasses non-human transgenic organisms, including animals and plants, that contains cells which express the transactivator fusion protein of the invention (i.e., a
15 nucleic acid encoding the transactivator is incorporated into one or more chromosomes in cells of the transgenic organism).

E. Homologous Recombinant Organisms

The invention also provides a homologous recombinant non-human organism
20 expressing the fusion protein of the invention. The term "homologous recombinant organism" as used herein is intended to describe an organism, e.g. animal or plant, containing a gene which has been modified by homologous recombination between the gene and a DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal. In one embodiment, the non-human animal is a mouse, although the invention is not limited thereto.
25 An animal can be created in which nucleic acid encoding the fusion protein has been introduced into a specific site of the genome, i.e., the nucleic acid has homologously recombined with an endogenous gene.

To create such a homologous recombinant animal, a vector is prepared which contains DNA encoding the fusion protein flanked at its 5' and 3' ends by additional nucleic acid of a
30 eukaryotic gene at which homologous recombination is to occur. The additional nucleic acid flanking that encoding the fusion protein is of sufficient length for successful homologous recombination with the eukaryotic gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see e.g., Thomas, K.R. and Capecchi, M. R. (1987) *Cell* 51:503 for a description of homologous recombination vectors). The vector is
35 introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected (see e.g., Li, E. et al. (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse) to form aggregation chimeras (see e.g., Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed.

(IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harbouring the homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA. These "germline
5 transmission" animals can further be mated to animals carrying a gene operatively linked to at least one *tet* operator sequence (discussed in more detail in Section III below).

In addition to the homologous recombination approaches described above, enzyme-assisted site-specific integration systems are known in the art and can be applied to the components of the regulatory system of the invention to integrate a DNA molecule at a
10 predetermined location in a second target DNA molecule. Examples of such enzyme-assisted integration systems include the Cre recombinase-lox target system (e.g., as described in Baubonis, W. and Sauer, B. (1993) *Nucl. Acids Res.* 21:2025-2029; and Fukushige, S. and Sauer, B. (1992) *Proc. Natl. Acad. Sci. USA* 89:7905-7909) and the FLP recombinase-FRT target system (e.g., as described in Dang, D.T. and Perrimon, N. (1992) *Dev. Genet.* 13:367-
15 375; and Fiering, S. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:8469-8473).

III. Target Transcription Units Regulated by a Tetracycline-Inducible Transactivator

A fusion protein of the invention is used to regulate the transcription of a target nucleotide sequence. This target nucleotide sequence is operatively linked to a regulatory
20 sequence to which the fusion protein binds. More specifically, the fusion protein regulates expression of a nucleotide sequence operatively linked to at least one *tet* operator sequence. Accordingly, another aspect of the invention relates to target nucleic acids (e.g., DNA molecules) comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence. Such nucleic acids are also referred to herein as *tet*-regulated
25 transcription units (or simply transcription units).

Within a transcription unit, the "nucleotide sequence to be transcribed" typically includes a minimal promoter sequence which is not itself transcribed but which serves (at least in part) to position the transcriptional machinery for transcription. The minimal promoter sequence is linked to the transcribed sequence in a 5' to 3' direction by
30 phosphodiester bonds (i.e., the promoter is located upstream of the transcribed sequence) to form a contiguous nucleotide sequence. Accordingly, as used herein, the terms "nucleotide sequence to be transcribed" or "target nucleotide sequence" are intended to include both the nucleotide sequence which is transcribed into mRNA and an operatively linked upstream minimal promoter sequence. The term "minimal promoter" is intended to describe a partial
35 promoter sequence which defines the start site of transcription for the linked sequence to be transcribed but which by itself is not capable of initiating transcription efficiently, if at all. Thus, the activity of such a minimal promoter is dependent upon the binding of a transcriptional activator (such as the tetracycline-inducible fusion protein of the invention) to an operatively linked regulatory sequence (such as one or more *tet* operator sequences). In

one embodiment, the minimal promoter is from the human cytomegalovirus (as described in Boshart et al. (1985) *Cell* 41:521-530). Preferably, nucleotide positions between about +75 to -53 and +75 to -31 are used. Other suitable minimal promoters are known in the art or can be identified by standard techniques. For example, a functional promoter which activates transcription of a contiguously linked reporter gene (e.g., chloramphenicol acetyl transferase, β -galactosidase or luciferase) can be progressively deleted until it no longer activates expression of the reporter gene alone but rather requires the presence of an additional regulatory sequence(s).

Within a transcription unit, the target nucleotide sequence (including the transcribed nucleotide sequence and its upstream minimal promoter sequence) is operatively linked to at least one *tet* operator sequence. In a typical configuration, the *tet* operator sequence(s) is operatively linked upstream (i.e., 5') of the minimal promoter sequence through a phosphodiester bond at a suitable distance to allow for transcription of the target nucleotide sequence upon binding of a regulatory protein (e.g., the transactivator fusion protein) to the *tet* operator sequence. That is, the transcription unit is comprised of, in a 5' to 3' direction: *tet* operator sequence(s) - a minimal promoter - a transcribed nucleotide sequence. It will be appreciated by those skilled in the art that there is some flexibility in the permissible distance between the *tet* operator sequence(s) and the minimal promoter, although typically the *tet* operator sequences will be located within about 200-400 base pairs upstream of the minimal promoter.

The nucleotide sequences of examples of *tet*-regulated promoters, containing *tet* operator sequences linked to a minimal promoter, that can be used in the invention are shown in SEQ ID NO: 8-10. The nucleotide sequences of SEQ ID NOs: 8 and 9 comprise a cytomegalovirus minimal promoter linked to ten *tet* operator sequences; the two nucleotide sequences differ in the distance between the operators and the first transcribed nucleotide. The nucleotide sequence of SEQ ID NO: 10 comprises a herpes simplex virus minimal tk promoter linked to ten *tet* operator sequences. The promoter of SEQ ID NO: 8 corresponds to P_hCMV*-1, described in Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551. The promoter of SEQ ID NO: 9 corresponds to P_hCMV*-2, also described in Gossen, M. and Bujard, H, *cited supra*.

Alternatively, since regulatory elements have been observed in the art to function downstream of sequences to be transcribed, it is likely that the *tet* operator sequence(s) can be operatively linked downstream (i.e., 3') of the transcribed nucleotide sequence. Thus, in this configuration, the transcription unit is comprised of, in a 5' to 3' direction: a minimal promoter - a transcribed nucleotide sequence - *tet* operator sequence(s). Again, it will be appreciated that there is likely to be some flexibility in the permissible distance downstream at which the *tet* operator sequence(s) can be linked.

The term "*tet* operator sequence" is intended to encompass all classes of *tet* operators (e.g., A, B, C, D and E). A nucleotide sequence to be transcribed can be operatively linked to

a single *tet* operator sequence, or for an enhanced range of regulation, it can be operatively linked to multiple *tet* operator sequences (e.g., two, three, four, five, six, seven, eight, nine, ten or more operator sequences). In a preferred embodiment, the sequence to be transcribed is operatively linked to seven *tet* operator sequences.

5 A *tet*-regulated transcription unit can further be incorporated into a recombinant vector (e.g., a plasmid or viral vector) by standard recombinant DNA techniques. The transcription unit, or recombinant vector in which it is contained, can be introduced into a host cell by standard transfection techniques, such as those described above. It should be appreciated that, after introduction of the transcription unit into a population of host cells, it
10 may be necessary to select a host cell clone which exhibit low basal expression of the *tet* operator-linked nucleotide sequence (i.e., selection for a host cell in which the transcription unit has integrated at a site that results in low basal expression of the *tet* operator-linked nucleotide sequence). Furthermore, a *tet*-regulated transcription unit can be introduced, by procedures described above, into the genome of a non-human animal at an embryonic stage
15 or into plant cells to create a transgenic or homologous recombinant organisms carrying the transcription unit in some or all of its cells. Again, it should be appreciated that it may be necessary to select a transgenic or homologous organism in which there is low basal expression of the the *tet* operator-linked nucleotide sequence in cells of interest.

 In one embodiment, the target nucleotide sequence of the *tet*-regulated transcription
20 unit encodes a protein of interest. Thus, upon induction of transcription of the nucleotide sequence by the transactivator of the invention and translation of the resultant mRNA, the protein of interest is produced in a host cell or animal. Alternatively, the nucleotide sequence to be transcribed can encode for an active RNA molecule, e.g., an antisense RNA molecule or ribozyme. Expression of active RNA molecules in a host cell or animal can be used to
25 regulate functions within the host (e.g., prevent the production of a protein of interest by inhibiting translation of the mRNA encoding the protein).

 A transactivator of the invention can be used to regulate transcription of an exogenous nucleotide sequence introduced into the host cell or animal. An "exogenous" nucleotide sequence is a nucleotide sequence which is introduced into the host cell and typically is
30 inserted into the genome of the host. The exogenous nucleotide sequence may not be present elsewhere in the genome of the host (e.g., a foreign nucleotide sequence) or may be an additional copy of a sequence which is present within the genome of the host but which is integrated at a different site in the genome. An exogenous nucleotide sequence to be transcribed and an operatively linked *tet* operator sequence(s) can be contained within a
35 single nucleic acid molecule which is introduced into the host cell or animal.

 Alternatively, a transactivator of the invention can be used to regulate transcription of an endogenous nucleotide sequence to which a *tet* operator sequence(s) has been linked. An "endogenous" nucleotide sequence is a nucleotide sequence which is present within the genome of the host. An endogenous gene can be operatively linked to a *tet* operator

sequence(s) by homologous recombination between a *tetO*-containing recombination vector and sequences of the endogeneous gene. For example, a homologous recombination vector can be prepared which includes at least one *tet* operator sequence and a minimal promoter sequence flanked at its 3' end by sequences representing the coding region of the endogenous gene and flanked at its 5' end by sequences from the upstream region of the endogenous gene by excluding the actual promoter region of the endogenous gene. The flanking sequences are of sufficient length for successful homologous recombination of the vector DNA with the endogenous gene. Preferably, several kilobases of flanking DNA are included in the homologous recombination vector. Upon homologous recombination between the vector DNA and the endogenous gene in a host cell, a region of the endogenous promoter is replaced by the vector DNA containing one or more *tet* operator sequences operably linked to a minimal promoter. Thus, expression of the endogenous gene is no longer under the control of its endogenous promoter but rather is placed under the control of the *tet* operator sequence(s) and the minimal promoter.

In another embodiment, *tet* operator sequences can be inserted elsewhere within an endogenous gene, preferably within a 5' or 3' regulatory region, via homologous recombination to create an endogenous gene whose expression can be regulated by a tetracycline-regulated fusion protein described herein. For example, one or more *tetO* sequences can be inserted into a promoter or enhancer region of an endogenous gene such that promoter or enhancer function is maintained (i.e., the *tetO* sequences are introduced into a site of the promoter/enhancer region that is not critical for promoter/enhancer function). Regions within promoters or enhancers which can be altered without loss of promoter/enhancer function are known in the art for many genes or can be determined by standard techniques for analyzing critical regulatory regions. An endogenous gene having *tetO* sequences inserted into a non-critical regulatory region will retain the ability to be expressed in its normal constitutive and/or tissue-specific manner but, additionally, can be downregulated by a tetracycline-controlled transcriptional inhibitor protein in a controlled manner. For example, constitutive expression of such a modified endogenous gene can be inhibited by in the presence of tetracycline (or analogue) using an inhibitor fusion protein that binds to *tetO* sequences in the presence of tetracycline (or analogue) (as described in further detail in Section IV and Section VI, Part B, below).

A. Regulation of Expression of tet Operator-Linked Nucleotide Sequences

Expression of a *tet* operator-linked nucleotide sequences is regulated by a transactivator fusion protein of the invention. Thus, the fusion protein and the target nucleic acid are both present in a host cell or organism. The presence of both the transactivator fusion protein and the target transcription unit in the same host cell or organism can be achieved in a number of different ways. For example, a host cell can be transfected with one nucleic acid of the expression system (e.g., encoding the transactivator fusion protein), stably

transfected cells can be selected and then the transfected cells can be re-transfected (also referred to as "supertransfected") with nucleic acid corresponding to the other nucleic acid of the expression system (e.g., the target nucleic acid to be transcribed). Two distinct selectable markers can be used for selection, e.g., uptake of the first nucleic acid can be selected with
5 G418 and uptake of the second nucleic acid can be selected with hygromycin. Alternatively, a single population of cells can be transfected with nucleic acid corresponding to both components of the system. Accordingly, the invention provides a nucleic acid composition comprising:

- 10 • a first nucleic acid encoding a fusion protein which activates transcription, the fusion protein comprising a first polypeptide which binds to a *tet* operator sequence in the presence of tetracycline or a tetracycline analogue operatively linked to a second polypeptide which activates transcription in eukaryotic cells; and
- 15 • a second nucleic acid comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.

In one embodiment, the two nucleic acids are two separate molecules (e.g., two different vectors). In this case, a host cell is cotransfected with the two nucleic acid molecules or
20 successively transfected first with one nucleic acid molecule and then the other nucleic acid molecule. In another embodiment, the two nucleic acids are linked (i.e., colinear) in the same molecule (e.g., a single vector). In this case, a host cell is transfected with the single nucleic acid molecule.

The host cell may be a cell cultured *in vitro* or a cell present *in vivo* (e.g., a cell
25 targeted for gene therapy). The host cell can further be a fertilized oocyte, embryonic stem cell or any other embryonic cell used in the creation of non-human transgenic or homologous recombinant animals. Transgenic or homologous recombinant animals which comprise both nucleic acid components of the expression system can be created by introducing both nucleic acids into the same cells at an embryonic stage, or more preferably, an animal which carries
30 one nucleic acid component of the system in its genome is mated to an animal which carries the other nucleic acid component of the system in its genome. Offspring which have inherited both nucleic acid components can then be identified by standard techniques.

B. Coordinate Regulation of Expression of Two Nucleotide Sequences

35 In addition to providing a system for the regulated expression of a single transcribed nucleotide sequence, the invention further permits coordinated regulation of the expression of two nucleotide sequences operatively linked to the same *tet* operator sequence(s). Accordingly, another aspect of the invention pertains to a novel *tet*-regulated transcription unit for coordinate regulation of two genes. In this transcription unit, the same *tet* operator

sequence(s) regulates the expression of two operatively linked nucleotide sequences that are transcribed in opposite directions from the common *tet* operator sequence(s). Accordingly, one nucleotide sequence is operatively linked to one side of the *tet* operator sequence (e.g., the 5' end on the top strand of DNA) and the other nucleotide sequence is operatively linked to the opposite side of the *tet* operator sequence (e.g., the 3' end on the top strand of DNA). Additionally, it should be understood that each nucleotide sequence to be transcribed includes an operatively linked minimal promoter sequence which is located between the nucleotide sequence to be transcribed and the *tet* operator sequence(s).

A representative example of such a transcription unit is diagrammed schematically in Figure 6. In this vectors, the two nucleotide sequences, operatively linked to the same *tet* operator sequence(s), are transcribed in opposite directions relative to the *tet* operator sequence(s) (i.e., the sequences are transcribed in a divergent manner upon activation by a transactivator fusion protein of the invention). By "transcribed in opposite directions relative to the *tet* operator sequence(s)", it is meant that the first nucleotide sequence is transcribed 5' to 3' from one strand of the DNA (e.g., the bottom strand) and the second nucleotide sequence is transcribed 5' to 3' from the other stand of the DNA (e.g., the top strand), resulting in bidirectional transcription away from the *tet* operator sequence(s).

Accordingly, the invention provides a recombinant vector for coordinately-regulated, bidirectional transcription of two nucleotide sequence. In one embodiment, the vector comprises a nucleotide sequence linked by phosphodiester bonds comprising, in a 5' to 3' direction:

- a first nucleotide sequence to be transcribed, operatively linked to
- at least one *tet* operator sequence, operatively linked to
- a second nucleotide sequence to be transcribed,

wherein transcription of the first and second nucleotide sequences proceeds in opposite directions from the at least one *tet* operator sequence(s) (i.e., the first and second nucleotide sequences are transcribed in a divergent manner).

In another embodiment, the vector does not include the first and second nucleotide sequence to be transcribed but instead contains cloning sites which allow for the introduction into the vector of nucleotide sequences of interest. Accordingly, in this embodiment, the vector comprises a nucleotide sequence comprising in a 5' to 3' direction:

- a first cloning site for introduction of a first nucleotide sequence to be transcribed, operatively linked to
- at least one *tet* operator sequence, operatively linked to
- a second cloning site for introduction of a second nucleotide sequence to be transcribed,

wherein transcription of a first and second nucleotide sequence introduced into the vector proceeds in opposite directions from the at least one *tet* operator sequence(s). It will be appreciated by those skilled in the art that this type of "cloning vector" may be in a form which also includes minimal promoter sequences such that a first nucleotide sequence introduced into the first cloning site is operatively linked to a first minimal promoter and a second nucleotide sequence introduced into the second cloning site is operatively linked to a second minimal promoter. Alternatively, the "cloning vector" may be in a form which does not include minimal promoter sequences and instead, nucleotide sequences including linked minimal promoter sequences are introduced into the cloning sites of the vector.

The term "cloning site" is intended to encompass at least one restriction endonuclease site. Typically, multiple different restriction endonuclease sites (e.g., a polylinker) are contained within the nucleic acid.

In yet another embodiment, the vector for coordinate, bidirectional transcription of two nucleotide sequences may contain a first nucleotide to be transcribed, such as that encoding a detectable marker (e.g., luciferase or β -galactosidase), and a cloning site for introduction of a second nucleotide sequence of interest.

The nucleotide sequences of two different suitable bidirectional promoter regions for use in a vector for coordinate regulation of two nucleotide sequences to be transcribed, as described herein, are shown in Figures 7A and 7B (SEQ ID NOS: 6 and 7, respectively). In the construct of Figure 7A, both minimal promoters present in the construct are derived from a CMV promoter. In the construct of Figure 7B, one minimal promoter present in the construct is derived from a CMV promoter, whereas the second minimal promoter is derived from a TK promoter. A plasmid pUHDG1316-8, comprising a bidirectional promoter of the invention, has been deposited on July 8, 1994 under the provisions of the Budapest Treaty at the Deutsche Sammlung Von Mikroorganismen und ZellKulturen GmbH (DSMZ) in Braunschweig, Germany and assigned deposit number DSM 9281.

The transcription unit of the invention for bidirectional transcription of two nucleotide sequences operatively linked to the same *tet* operator sequence(s) is useful for coordinating the expression of the two nucleotide sequences of interest. Preferably, at least one of the nucleotide sequences to be transcribed is a eukaryotic nucleotide sequence. In one application, the vector is used to produce stoichiometric amounts of two subunits of a heterodimeric molecule in the same cell. For example, the vector can be used produce antibody heavy and light chains in the same cell or to produce growth factor receptor subunits in the same cells. In another application, the vector is used to express two gene products that cooperate in establishing a particular cellular phenotype. In yet another application, the vector is used to coexpress an indicator function and a gene of interest, wherein the indicator is utilized to monitor expression of the gene of interest. Thus, one of the two coordinately expressed sequences can encode a gene of interest and the other can encode a detectable

marker, such as a surface marker or enzyme (e.g., β -galactosidase or luciferase) which is used for selection of cells expressing the gene of interest.

Transcription of the two coordinately-regulated nucleotide sequences can be induced by tetracycline (or an analogue thereof) by use of the Tc-inducible transcriptional activator of the invention to regulate expression of the two nucleotide sequences. Thus, in this system, expression of both nucleotide sequences is "off" in the absence of Tc (or analogue), whereas expression is turned "on" by the presence of Tc (or analogue). Alternatively, the vector for coordinate regulation of two nucleotide sequences can be used in conjunction with other tetracycline-regulated transcription factors known in the art. For example, a transactivator fusion protein of a wild-type Tet repressor fused to a transcriptional activation domain, which activates gene expression in the absence of Tc (or analogue), such as the tTA described in Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551, can also be used in conjunction with this target transcription unit for coordinate regulation.

15 C. Independent Regulation of Expression of Multiple Nucleotide Sequences

The invention still further permits independent and opposite regulation of two or more nucleotide sequences to be transcribed. Accordingly, another aspect of the invention pertains to a novel *tet*-regulated transcription unit for independent regulation of two or more genes. To independently regulate the expression of two nucleotide sequences to be transcribed, one nucleotide sequence is operatively linked to a *tet* operator sequence(s) of one class type and the other nucleotide sequence is operatively linked to a *tet* operator sequence(s) of another class type. Accordingly, the invention provides at least one recombinant vector for independent regulation of transcription of two nucleotide sequences. In one embodiment, the vector(s) comprises:

- a first nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence of a first class type; and
- a second nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence of a second class type.

(It should be understood that each nucleotide sequence to be transcribed also includes an operatively linked, upstream minimal promoter sequence.) The two independently regulated transcription units can be included on a single vector, or alternatively, on two separate vectors. The recombinant vector(s) containing the nucleotide sequences to be transcribed can be introduced into a host cell or animal as described previously.

In another embodiment, the vector(s) does not include the first and second nucleotide sequence to be transcribed but instead contains cloning sites which allow for the introduction into the vector of nucleotide sequences of interest. Accordingly, in this embodiment, the vector(s) comprises:

- a first cloning site for introduction of a first nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence of a first class type; and
- 5 • a second cloning site for introduction of a second nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence of a second class type.

This cloning vector(s) may be in a form that already includes first and second minimal promoters operatively linked, respectively, to the first and second cloning sites.

- 10 Alternatively, nucleotide sequences to be transcribed which include an operatively linked minimal promoter can be introduced into the cloning vector.

In yet another embodiment, the vector for independent regulation of two nucleotide sequences may contain a first nucleotide to be transcribed, such as that encoding a detectable marker or a suicide gene, operatively linked to at least one *tet* operator sequence of a first class type and a cloning site for introduction of a second nucleotide sequence of interest such that it is operatively linked to at least one *tet* operator sequence of a second class type.

It will be appreciated by those skilled in the art that various combinations of classes of *tet* operator sequences can be used for independent regulation of two nucleotide sequences. For example, the first *tet* operator sequence(s) can be of the class A type and the second can be of the class B type, or the first *tet* operator sequence can be of the class B type and the second can be of the class C type, etc. Preferably, one to the two *tet* operators used is a class B type operator.

Independent transcription of the first and second nucleotide sequences is regulated in a host cell by further introducing into the host cell one or more nucleic acids encoding two different transactivator fusion proteins which bind independently to *tet* operator sequences of different class types. The first fusion protein comprises a polypeptide which binds to a *tet* operator sequence in the presence of tetracycline or a tetracycline analogue, operatively linked to a polypeptide which activates transcription in eukaryotic cells (e.g., a transactivator fusion protein of the invention, such as a mutated Tn10-derived Tet repressor linked to a VP16 activation region). The second fusion protein comprises a polypeptide which binds to a *tet* operator sequence in the absence of tetracycline or a tetracycline analogue, operatively linked to a polypeptide which activates transcription in eukaryotic cells (e.g., a wild-type Tn10-derived Tet repressor linked to a VP16 activation region, such as the tTA described in Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551). In one embodiment, the first fusion protein binds to the *tet* operator sequence of the first class type used in the transcription unit and the second fusion protein binds to the *tet* operator sequence of the second class type used in the transcription unit. Alternatively, in another embodiment, the first fusion protein binds to the second class type of *tet* operator and the second fusion protein binds to the first class type of *tet* operator.

For example, the first nucleotide sequence to be transcribed may be linked to a class A *tet* operator and the first fusion protein may bind to class A operators, whereas the second nucleotide sequence to be transcribed may be linked to a class B *tet* operator and the second fusion protein may bind to class B operators. Thus, in this embodiment, transcription of the first nucleotide sequence is activated in the presence of Tc (or analogue thereof) while transcription of the second nucleotide sequence is activated in the absence of Tc (or analogue thereof). Alternatively, in another embodiment, the first fusion protein binds to class B operators and the second fusion protein binds to class A operators. In this case, transcription of the second nucleotide sequence is activated in the presence of Tc (or analogue thereof) while transcription of the first nucleotide sequence is activated in the absence of Tc (or analogue thereof). Appropriate transactivator proteins for use in this system can be designed as described above in Section I and in Gossen and Bujard (1992) cited *supra*. In order to inhibit heterodimerization between the two different types of Tet repressor fusion proteins present in the same cell, it may be necessary to mutate the dimerization region of one or both of the transactivator fusion proteins. Mutations can be targeted to the C-terminal region of TetR known to be involved in dimerization. The dimerization region has been described in detail based upon the crystal structure of TetR (see Hinrichs, W. et al. (1994) *Science* 264:418-420).

This system allows for independent and opposite regulation of the expression of two genes by Tc and analogues thereof. Use of different Tc analogues as inducing agents may further allow for high, low or intermediate levels of expression of the different sequences (discussed in greater detail in Section V below). The novel transcription unit of the invention for independently regulating the expression of two genes, described above, can be used in situations where two gene products are to be expressed in the same cell but where it is desirable to express one gene product while expression of the other gene product is turned "off", and vice versa. For example, this system is particularly useful for expressing in the same host cell either a therapeutic gene or a suicide gene (i.e., a gene which encodes a product that can be used to destroy the cell, such as ricin or herpes simplex virus thymidine kinase). In many gene therapy situations, it is desirable to be able to express a gene for therapeutic purposes in a host cell but also to have the capacity to destroy the host cell once the therapy is completed. This can be accomplished using the above-described system by linking the therapeutic gene to one class of *tet* operator and the suicide gene to another class of *tet* operator. Thus, expression of the therapeutic gene in a host cell can be stimulated by Tc (in which case expression of the suicide gene is absent). Then, once the therapy is complete, Tc is removed, which turns off expression of the therapeutic gene and turns on expression of the suicide gene in the cell.

D. Combined Coordinate and Independent Regulation of Multiple Nucleotide Sequences

It is further possible to regulate the expression of four nucleotide sequences by combining the system described in Section IIIB with the system described in Section IIIC such that two pairs of sequences are coordinately regulated while one pair is independently regulated from the other pair. Accordingly, two target transcription units can be designed comprising:

- a first nucleic acid comprising in a 5' to 3' direction: a first nucleotide sequence to be transcribed, a *tet* operator sequence(s) of a first class type, and a second nucleotide sequence to be transcribed
- a second nucleic acid comprising in a 5' to 3' direction: a third nucleotide sequence to be transcribed, a *tet* operator sequence(s) of a second class type, and a fourth nucleotide sequence to be transcribed.

Transcription of the first and second nucleotide sequences in the first nucleic acid proceeds in a divergent manner from the first class of *tet* operator sequence(s). Likewise, transcription of the third and fourth nucleotide sequences in the second nucleic acid proceeds in a divergent manner from the second class of *tet* operator sequence(s). Thus, expression of the first and second nucleotide sequences is coordinately regulated and expression of the third and fourth nucleotide sequences is coordinately regulated. However, expression of the first and second sequences is independently (and oppositely) regulated compared to the third and fourth sequences through the use of two different transactivator fusion proteins, as described above, one which activates transcription in the presence of Tc (or analogue thereof) and the other which activates transcription in the absence of Tc (or analogue thereof). One transactivator is designed to bind to a *tet* operators of the first class type and the other is designed to bind to a *tet* operators of the second class type. In other embodiments, rather than already containing first, second, third and/or fourth nucleotide sequences to be transcribed, these transcription units can contain cloning sites which allow for the introduction of first, second, third and/or fourth nucleotide sequences to be transcribed.

IV. Tetracycline-Regulated Transcriptional Inhibitors

Another aspect of the invention pertains to transcriptional inhibitor fusion proteins. The inhibitor fusion proteins of the invention are constructed similarly to the transactivator fusion proteins of the invention (see Section I above) but instead of containing a polypeptide domain that stimulates transcription in eukaryotic cells, the inhibitor fusion proteins contain a polypeptide domain that inhibits transcription in eukaryotic cells. The inhibitor fusion proteins are used to downregulate the expression of genes operably linked to *tetO* sequences. For example, when a *tetO*-linked gene is introduced into a host cell or animal, the level of

basal, constitutive expression of the gene may vary depending upon the type of cell or tissue in which the gene is introduced and on the site of integration of the gene. Alternatively, constitutive expression of endogenous genes into which *tetO* sequences have been introduced may vary depending upon the strength of additional endogenous regulatory sequences in the vicinity. The inhibitor fusion proteins described herein provide compositions that can be used to inhibit the expression of such *tetO*-linked genes in a controlled manner.

In one embodiment, the inhibitor fusion protein of the invention comprises a first polypeptide that binds to *tet* operator sequences in the absence, but not the presence, of tetracycline (Tc) or an analogue thereof operatively linked to a heterologous second polypeptide that inhibits transcription in eukaryotic cells. In another embodiment, the inhibitor fusion protein comprises a first polypeptide that binds to *tet* operator sequences in the presence, but not the absence, of tetracycline operatively linked to a heterologous second polypeptide that inhibits transcription in eukaryotic cells. The term "heterologous" is intended to mean that the second polypeptide is derived from a different protein than the first polypeptide. Like the transactivator fusion proteins, the transcriptional inhibitor fusion proteins can be prepared using standard recombinant DNA techniques as described herein.

A. The first polypeptide of the transcriptional inhibitor fusion protein

The transcriptional inhibitor fusion protein of the invention is composed, in part, of a first polypeptide which binds to a *tet* operator sequence either (i) in the absence, but not the presence of tetracycline (Tc), or an analogue thereof, or alternatively, (ii) in the presence, but not the absence of Tc or an analogue thereof.

Preferably, in the former embodiment, the first polypeptide is a wild-type Tet repressor (which binds to *tet* operator sequences in the absence but not the presence of Tc). A wild-type Tet repressor of any class (e.g., A, B, C, D or E) may be used as the first polypeptide. Preferably, the wild-type Tet repressor is a Tn10-derived Tet repressor. The nucleotide and amino acid sequences of a wild-type Tn10-derived Tet repressor are shown in SEQ ID NO: 16 and SEQ ID NO: 17, respectively.

Alternatively, in the latter embodiment, the first polypeptide is a mutated Tet repressor as described in Section I, part A above (which binds to *tet* operator sequences in the presence but not the absence of Tc). A mutated Tet repressor of any class (e.g., A, B, C, D or E) may be used as the first polypeptide. Preferably, the mutated Tet repressor is a Tn10-derived Tet repressor having one or more amino acid substitutions at positions 71, 95, 101 and/or 102. The nucleotide and amino acid sequences of such a mutated Tn10-derived Tet repressor are shown in SEQ ID NO: 18 and SEQ ID NO: 19, respectively.

B. The second polypeptide of the transcriptional inhibitor fusion protein

The first polypeptide of the transcriptional inhibitor fusion protein is operatively linked to a second polypeptide which directly or indirectly inhibits transcription in eukaryotic

cells. As described in Section I, above, to operatively link the first and second polypeptides of a fusion protein, typically nucleotide sequences encoding the first and second polypeptides are ligated to each other in-frame to create a chimeric gene encoding the fusion protein.

However, the first and second polypeptides can be operatively linked by other means that preserve the function of each polypeptide (e.g., chemically crosslinked). Although the fusion proteins are typically described herein as having the first polypeptide at the amino-terminal end of the fusion protein and the second polypeptide at the carboxy-terminal end of the fusion protein, it will be appreciated by those skilled in the art that the opposite orientation (i.e., the second polypeptide at the amino-terminal end and the first polypeptide at the carboxy-terminal end) is also contemplated by the invention.

Proteins and polypeptide domains within proteins which can function to inhibit transcription in eukaryotic cells have been described in the art (for reviews see, e.g., Renkawitz, R. (1990) *Trends in Genetics* 6:192-197; and Herschbach, B.M. and Johnson, A.D. (1993) *Annu. Rev. Cell. Biol.* 2:479-509). Such transcriptional inhibitor domains have been referred to in the art as "silencing domains" or "repressor domains." Although the precise mechanism by which many of these polypeptide domains inhibit transcription is not known (and the invention is not intended to be limited by mechanism), there are several possible means by which repressor domains may inhibit transcription, including: 1) competitive inhibition of binding of either activator proteins or the general transcriptional machinery, 2) prevention of the activity of a DNA bound activator and 3) negative interference with the assembly of a functional preinitiation complex of the general transcription machinery. Thus, a repressor domain may have a direct inhibitory effect on the transcriptional machinery or may inhibit transcription indirectly by inhibiting the activity of activator proteins. Accordingly, the term "a polypeptide that inhibits transcription in eukaryotic cells" as used herein is intended to include polypeptides which act either directly or indirectly to inhibit transcription. As used herein, "inhibition" of transcription is intended to mean a diminution in the level or amount of transcription of a target gene compared to the level or amount of transcription prior to regulation by the transcriptional inhibitor protein. Transcriptional inhibition may be partial or complete. The terms "silencer", "repressor" and "inhibitor" are used interchangeably herein to describe a regulatory protein, or domains thereof, that can inhibit transcription.

A transcriptional "repressor" or "silencer" domain as described herein is a polypeptide domain that retains its transcriptional repressor function when the domain is transferred to a heterologous protein. Proteins which have been demonstrated to have repressor domains that can function when transferred to a heterologous protein include the v-erbA oncogene product (Baniahmad, A. et al. (1992) *EMBO J.* 11:1015-1023), the thyroid hormone receptor (Baniahmad, *supra*), the retinoic acid receptor (Baniahmad, *supra*), and the Drosophila Krueppel (Kr) protein (Licht, J.D. et al. (1990) *Nature* 346:76-79; Sauer, F. and Jäckle, H. (1991) *Nature* 353:563-566; Licht, J.D. et al. (1994) *Mol. Cell. Biol.* 14:4057-4066). Non-

limiting examples of other proteins which have transcriptional repressor activity in eukaryotic cells include the *Drosophila* homeodomain protein even-skipped (*eve*), the *S. cerevisiae* Ssn6/Tup1 protein complex (see Herschbach and Johnson, *supra*), the yeast SIR1 protein (see Chien, et al. (1993) *Cell* 75:531-541), NeP1 (see Kohne, et al. (1993) *J. Mol. Biol.* 232:747-755), the *Drosophila* dorsal protein (see Kirov, et al. (1994) *Mol. Cell. Biol.* 14:713-722; Jiang, et al. (1993) *EMBO J.* 12:3201-3209), TSF3 (see Chen, et al. (1993) *Mol. Cell. Biol.* 13:831-840), SF1 (see Targa, et al. (1992) *Biochem. Biophys. Res. Comm.* 188:416-423), the *Drosophila* hunchback protein (see Zhang, et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:7511-7515), the *Drosophila* knirps protein (see Gerwin, et al. (1994) *Mol. Cell. Biol.* 14:7899-7908), the WT1 protein (Wilm's tumor gene product) (see Anant, et al. (1994) *Oncogene* 2:3113-3126; Madden et al., (1993) *Oncogene* 8:1713-1720), Oct-2.1 (see Lillycrop, et al. (1994) *Mol. Cell. Biol.* 14:7633-7642), the *Drosophila* engrailed protein (see Badiani, et al. (1994) *Genes Dev.* 8:770-782; Han and Manley, (1993) *EMBO J.* 12:2723-2733), E4BP4 (see Cowell and Hurst, (1994) *Nucleic Acids Res.* 22:59-65) and ZF5 (see Numoto, et al. (1993) *Nucleic Acids Res.* 21:3767-3775),

In a preferred embodiment, the second polypeptide of the transcriptional inhibitor fusion protein of the invention is a transcriptional silencer domain of the *Drosophila* Krueppel protein. A C-terminal region having repressor activity can be used, such as amino acids 403-466 of the native protein (see Sauer, F. and Jäckle, H., *supra*). This region is referred to as C64KR. The nucleotide and amino acid sequences of C64KR are shown in SEQ ID NO: 20 and SEQ ID NO: 21, respectively. Construction of an expression vector encoding a TetR-C64KR fusion protein is described in Example 4. Alternatively, an alanine-rich amino terminal region of Kr that also has repressor activity can be used as the second polypeptide of the fusion protein. For example, amino acids 26-110 of Kr (see Licht, J.D. et al., (1990) *supra*) can be used as the second polypeptide. Alternatively, shorter or longer polypeptide fragments encompassing either of the Kr silencer domains that still retain full or partial inhibitor activity are also contemplated (e.g., amino acids 62 to 92 of the N-terminal silencer domain; see Licht, et al. (1994) *supra*).

In another preferred embodiment, the second polypeptide of the transcriptional inhibitor fusion protein of the invention is a transcriptional silencer domain of the v-erbA oncogene product. The silencer domain of v-erbA has been mapped to approximately amino acid residues 362-632 of the native v-erbA oncogene product (see Baniahmad, et al. *supra*). Accordingly, a fragment encompassing this region is used as the second polypeptide of the silencer domain. In one embodiment, amino acid residues 364-635 of the native v-erbA protein are used. The nucleotide and amino acid sequences of this region of v-erbA are shown in SEQ ID NO: 22 and SEQ ID NO: 23, respectively. Construction of an expression vector encoding a TetR-v-erbA fusion protein is described in Example 5. Alternatively, shorter or longer polypeptide fragments encompassing the v-erbA silencer region that still retain full or partial inhibitor activity are also contemplated. For example, a.a. residues 346-

639, 362-639, 346-632, 346-616 and 362-616 of v-erbA may be used. Additionally, polypeptide fragments encompassing these regions that have internal deletions yet still retain full or partial inhibitor activity are encompassed by the invention, such as a.a. residues 362-468/508-639 of v-erbA. Furthermore, two or more copies of the silencer domain may be included in the fusion protein, such as two copies of a.a. residues 362-616 of v-erbA. Suitable silencer polypeptide domains of v-erbA are described further in Baniahmad, A. et al. (*supra*).

In other embodiments, other silencer domains are used. Non-limiting examples of polypeptide domains that can be used include: amino acid residues 120-410 of the thyroid hormone receptor alpha (THR α), amino acid residues 143-403 of the retinoic acid receptor alpha (RAR α), amino acid residues 186-232 of knirps, the N-terminal region of WT 1 (see Anant, *supra*), the N-terminal region of Oct-2.1 (see Lillycrop, *supra*), a 65 amino acid domain of E4BP4 (see Cowell and Hurst, *supra*) and the N-terminal zinc finger domain of ZF5 (see Numoto, *supra*). Moreover, shorter or longer polypeptide fragments encompassing these regions that still retain full or partial inhibitor activity are also contemplated.

In addition to previously described transcriptional inhibitor domains, novel transcriptional inhibitor domains, which can be identified by standard techniques, are within the scope of the invention. The transcriptional inhibitor ability of a polypeptide can be assayed by: 1) constructing an expression vector that encodes the test silencer polypeptide linked to another polypeptide having DNA binding activity (i.e., constructing a DNA binding domain-silencer domain fusion protein), 2) cotransfecting this expression vector into host cells together with a reporter gene construct that is normally constitutively expressed in the host cell and also contains binding sites for the DNA binding domain and 3) determining the amount of transcription of the reporter gene construct that is inhibited by expression of the fusion protein in the host cell. For example, a standard assay used in the art utilizes a fusion protein of a GAL4 DNA binding domain (e.g., amino acid residues 1-147) and a test silencer domain. This fusion protein is then used to inhibit expression of a reporter gene construct that contains positive regulatory sequences (that normally stimulate constitutive transcription) and GAL4 binding sites (see e.g., Baniahmad, *supra*).

C. A third polypeptide of the transcriptional inhibitor fusion protein

In addition to a Tet repressor and a transcriptional silencer domain, a transcriptional inhibitor fusion protein of the invention can contain an operatively linked third polypeptide which promotes transport of the fusion protein to a cell nucleus. As described for the transactivator fusion proteins (see Section I, Part C, above), a nuclear localization signal can be incorporated into the transcriptional inhibitor fusion protein.

D. Expression of the transcriptional inhibitor fusion protein

A nucleic acid molecule encoding a transcriptional inhibitor fusion protein of the invention can be incorporated into a recombinant expression vector and introduced into a host cell to express the fusion protein in the host cell as described in Section II, Parts A, B and C, above. Preferably, a host cell expressing a transcriptional inhibitor fusion protein of the invention also carries a *tet* operator-linked gene of interest (i.e., target nucleotide sequence to be transcribed).

Transgenic organisms expressing a transcriptional inhibitor fusion protein in cells thereof can be prepared as described in Section II, Part D, above. Moreover, homologous recombinant organisms expressing a transcriptional inhibitor fusion protein in cells thereof are also encompassed by the invention and can be prepared as described in Section II, Part E, above. The invention provides recombinant expression vectors suitable for homologous recombination. In one embodiment, such an expression vector comprises a nucleic acid molecule encoding a transcriptional inhibitor fusion protein of the invention which is flanked at its 5' and 3' ends by additional nucleic acid of a eukaryotic gene, the additional nucleic acid being of sufficient length for successful homologous recombination with the eukaryotic gene. Vectors and methods for creating homologous recombinant organisms that express the components of the regulatory system of the invention, and uses therefor, are described in further detail in U.S. Patent Application Serial Number 08/260,452. Preferably, a transgenic or homologous recombinant organism of the invention expressing a transcriptional inhibitor fusion protein in cells thereof also carries a *tet* operator-linked gene of interest (i.e., target nucleotide sequence to be transcribed) in cells thereof.

V. Kits of the Invention

Another aspect of the invention pertains to kits which include the components of the inducible regulatory system of the invention. Such a kit can be used to regulate the expression of a gene of interest (i.e., a nucleotide sequence of interest to be transcribed) which can be cloned into a target transcription unit. The kit may include nucleic acid encoding a transcriptional activator fusion protein or a transcriptional inhibitor fusion protein or both. Alternatively, eukaryotic cells which have nucleic acid encoding a transactivator and/or inhibitor fusion protein stably incorporated therein, such that the transactivator and/or inhibitor fusion protein are expressed in the eukaryotic cell, may be provided in the kit.

In one embodiment, the kit includes a carrier means having in close confinement therein at least two container means: a first container means which contains a first nucleic acid (e.g., DNA) encoding a transactivator fusion protein of the invention (e.g., a recombinant expression vector encoding a first polypeptide which binds to a *tet* operator sequence in the presence of tetracycline operatively linked to a second polypeptide which activates transcription in eukaryotic cells), and a second container means which contains a second target nucleic acid (e.g., DNA) for the transactivator into which a nucleotide sequence of

interest can be cloned. The second nucleic acid typically comprises a cloning site for introduction of a nucleotide sequence to be transcribed (optionally including an operatively linked minimal promoter sequence) and at least one operatively linked *tet* operator sequence. The term "cloning site" is intended to encompass at least one restriction endonuclease site.

- 5 Typically, multiple different restriction endonuclease sites (e.g., a polylinker) are contained within the nucleic acid.

To regulate expression of a nucleotide sequence of interest using the components of the kit, the nucleotide sequence is cloned into the cloning site of the target vector of the kit by conventional recombinant DNA techniques and then the first and second nucleic acids are
10 introduced into a host cell or animal. The transactivator fusion protein expressed in the host cell or animal then regulates transcription of the nucleotide sequence of interest in the presence of the inducing agent (Tc or analogue thereof).

Alternatively, in another embodiment, the kit includes a eukaryotic cell which is stably transfected with a nucleic acid encoding a transactivator fusion protein of the invention
15 such that the transactivator is expressed in the cell. Thus, rather than containing nucleic acid alone, the first container means described above can contain a eukaryotic cell line into which the first nucleic acid encoding the transactivator has been stably introduced (e.g., by stable transfection by a conventional method such as calcium phosphate precipitation or electroporation, etc.). In this embodiment, a nucleotide sequence of interest is cloned into the
20 cloning site of the target vector of the kit and then the target vector is introduced into the eukaryotic cell expressing the transactivator fusion protein.

Alternatively or additionally, a recombinant vector of the invention for coordinate regulation of expression of two nucleotide sequences can also be incorporated into a kit of the invention. The vector can be included in the kit in a form that allows for introduction into the
25 vector of two nucleotide sequences of interest. Thus, in another embodiment, a kit of the invention includes 1) a first nucleic acid encoding a transactivator fusion protein of the invention (or a eukaryotic cell into which the nucleic acid has been stably introduced) and 2) a second nucleic acid comprising a nucleotide sequence comprising in a 5' to 3' direction: a first cloning site for introduction of a first nucleotide sequence of interest operatively linked
30 to at least one *tet* operator sequence operatively linked to a second cloning site for introduction of a second nucleotide sequence of interest, wherein transcription of the first and second nucleotide sequences proceeds in opposite directions from the at least one *tet* operator sequence. Optionally, the vector can include operatively linked minimal promoter sequences. In another embodiment, the vector can be in a form that already contains one nucleotide
35 sequence to be transcribed (e.g., encoding a detectable marker such as luciferase, β -galactosidase or CAT) and a cloning site for introduction of a second nucleotide sequence of interest to be transcribed.

The transcription units and transactivators of the invention for independent regulation of expression of two nucleotide sequences to be transcribed can also be incorporated into a kit

of the invention. The target transcription units can be in a form which allows for introduction into the transcription units of nucleotide sequences of interest to be transcribed. Thus, in another embodiment, a kit of the invention includes 1) a first nucleic acid encoding a transactivator which binds to a *tet* operator of a first class type in the presence of Tc or an analogue thereof, 2) a second nucleic acid comprising a first cloning site for introduction of a first nucleotide sequence to be transcribed operatively linked to at least one *tet* operator of a first class type, 3) a third nucleic acid encoding a transactivator which binds to a *tet* operator of a second class type in the absence of Tc or an analogue thereof, , and 4) a fourth nucleic acid comprising a second cloning site for introduction of a second nucleotide sequence to be transcribed operatively linked to at least one *tet* operator of a second class type. (Optionally, minimal promoter sequences are included in the second and fourth nucleic acids). In another embodiment, one nucleotide sequence to be transcribed (e.g., encoding a suicide gene) is already contained in either the second or the fourth nucleic acid. In yet another embodiment, the nucleic acids encoding the transactivators (e.g., the first and third nucleic acids described above) can be stably introduced into a eukaryotic cell line which is provided in the kit.

In yet another embodiment, a kit of the invention includes a first container means containing a first nucleic acid encoding a transcriptional inhibitor fusion protein of the invention (e.g., the fusion protein inhibits transcription in eukaryotic cells either only in the presence of Tc or only in the absence of Tc) and a second container means containing a second nucleic acid comprising a cloning site for introduction of a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence. The kit may further include a third nucleic acid encoding a transactivator fusion protein that binds to *tetO* sequences either only in the presence of Tc or only in the absence of Tc. Alternatively, the first and/or third nucleic acids (i.e., encoding the inhibitor or transactivator fusion proteins) may be stably incorporated into a eukaryotic host cell which is provided in the kit.

In still another embodiment, a kit of the invention may include at least one tetracycline or tetracycline analogue. For example, the kit may include a container means which contains tetracycline, anhydrotetracycline, doxycycline, epioxytetracycline or other tetracycline analogue described herein.

30

VI. Regulation of Gene Expression by Tetracycline or Analogues Thereof

A. Stimulation of Gene Expression by Transactivator Fusion Proteins

In a host cell which carries nucleic acid encoding a transactivator fusion protein of the invention and a nucleotide sequence operatively linked to the *tet* operator sequence (i.e., gene of interest to be transcribed), high level transcription of the nucleotide sequence operatively linked to the *tet* operator sequence(s) does not occur in the absence of the inducing agent, tetracycline or analogues thereof. The level of basal transcription of the nucleotide sequence may vary depending upon the host cell and site of integration of the sequence, but is

35

generally quite low or even undetectable in the absence of Tc. In order to induce transcription in a host cell, the host cell is contacted with tetracycline or a tetracycline analogue. Accordingly, another aspect of the invention pertains to methods for stimulating transcription of a nucleotide sequence operatively linked to a *tet* operator sequence in a host cell or animal which expresses a transactivator fusion protein of the invention. The methods involve contacting the cell with tetracycline or a tetracycline analogue or administering tetracycline or a tetracycline analogue to a subject containing the cell.

The term "tetracycline analogue" is intended to include compounds which are structurally related to tetracycline and which bind to the Tet repressor with a K_a of at least about 10^6 M^{-1} . Preferably, the tetracycline analogue binds with an affinity of about 10^9 M^{-1} or greater. Examples of such tetracycline analogues include, but are not limited to, anhydrotetracycline, doxycycline, chlorotetracycline, oxytetracycline and others disclosed by Hlavka and Boothe, "The Tetracyclines," in *Handbook of Experimental Pharmacology* 78, R.K. Blackwood et al. (eds.), Springer-Verlag, Berlin-New York, 1985; L.A. Mitscher, "The Chemistry of the Tetracycline Antibiotics", *Medicinal Research* 9, Dekker, New York, 1978; Noyee Development Corporation, "Tetracycline Manufacturing Processes" *Chemical Process Reviews*, Park Ridge, NJ, 2 volumes, 1969; R.C. Evans, "The Technology of the Tetracyclines", *Biochemical Reference Series 1*, Quadrangle Press, New York, 1968; and H.F. Dowling, "Tetracycline", *Antibiotic Monographs*, no. 3, Medical Encyclopedia, New York, 1955. Preferred Tc analogues for high level stimulation of transcription are anhydrotetracycline and doxycycline. A Tc analogue can be chosen which has reduced antibiotic activity compared to Tc. Examples of such Tc analogues are anhydrotetracycline, epioxytetracycline and cyanotetracycline.

To induce gene expression in a cell *in vitro*, the cell is contacted with Tc or a Tc analogue by culturing the cell in a medium containing the compound. When culturing cells *in vitro* in the presence of Tc or Tc analogue, a preferred concentration range for the inducing agent is between about 10 and about 1000 ng/ml. Tc or a Tc analogue can be directly added to media in which cells are already being cultured, or more preferably for high levels of gene induction, cells are harvested from Tc-free media and cultured in fresh media containing Tc, or an analogue thereof.

To induce gene expression *in vivo*, cells within in a subject are contacted with Tc or a Tc analogue by administering the compound to the subject. The term "subject" is intended to include humans and other non-human mammals including monkeys, cows, goats, sheep, dogs, cats, rabbits, rats, mice, and transgenic and homologous recombinant species thereof. Furthermore, the term "subject" is intended to include plants, such as transgenic plants. When the inducing agent is administered to a human or animal subject, the dosage is adjusted to preferably achieve a serum concentration between about 0.05 and 1.0 $\mu\text{g/ml}$. Tc or a Tc analogue can be administered to a subject by any means effective for achieving an *in vivo* concentration sufficient for gene induction. Examples of suitable modes of administration

include oral administration (e.g., dissolving the inducing agent in the drinking water), slow release pellets and implantation of a diffusion pump. To administer Tc or a Tc analogue to a transgenic plant, the inducing agent can be dissolved in water administered to the plant.

The ability to use different Tc analogues as inducing agents in this system allows for
5 modulate the level of expression of a *tet* operator-linked nucleotide sequence. As demonstrated in Example 2, anhydrotetracycline and doxycycline have been found to be strong inducing agents. The increase in transcription of the target sequence is typically as high as 1000- to 2000-fold, and induction factors as high as 20,000 fold can be achieved. Tetracycline, chlorotetracycline and oxytetracycline have been found to be weaker inducing
10 agents, i.e., the increase in transcription of a target sequence is in the range of about 10-fold. Thus, an appropriate tetracycline analogue is chosen as an inducing agent based upon the desired level of induction of gene expression. It is also possible to change the level of gene expression in a host cell or animal over time by changing the Tc analogue used as the inducing agent. For example, there may be situations where it is desirable to have a strong
15 burst of gene expression initially and then have a sustained lower level of gene expression. Accordingly, an analogue which stimulates a high levels of transcription can be used initially as the inducing agent and then the inducing agent can be switched to an analogue which stimulates a lower level of transcription. Moreover, when regulating the expression of multiple nucleotide sequences (e.g., when one sequence is regulated by a one of class *tet*
20 operator sequence(s) and the other is regulated by another class of *tet* operator sequence(s), as described above in Section III, Part C, above), it may be possible to independently vary the level of expression of each sequence depending upon which transactivator fusion protein is used to regulate transcription and which Tc analogue(s) is used as the inducing agent. Different transactivator fusion proteins are likely to exhibit different levels of responsiveness
25 to Tc analogues. The level of induction of gene expression by a particular combination of transactivator fusion protein and inducing agent (Tc or Tc analogue) can be determined by techniques described herein, (e.g., see Example 2). Additionally, the level of gene expression can be modulated by varying the concentration of the inducing agent. Thus, the expression system of the invention provides a mechanism not only for turning gene expression on or off,
30 but also for "fine tuning" the level of gene expression at intermediate levels depending upon the type and concentration of inducing agent used.

B. Inhibition of Gene Expression by Transcriptional Inhibitor Fusion Proteins

The invention also provides methods for inhibiting gene expression using the
35 transcriptional inhibitor fusion proteins of the invention. These methods can be used to down-regulate basal, constitutive or tissue-specific transcription of a *tetO*-linked gene of interest. For example, a gene of interest that is operatively linked to *tetO* sequences and additional positive regulatory elements (e.g., constitutive or tissue-specific enhancer sequences) will be transcribed in host cells at a level that is primarily determined by the

strength of the positive regulatory elements in the host cell. Moreover, a gene of interest that is operatively linked to *tetO* sequences and only a minimal promoter sequence may exhibit varying degrees of basal level transcription depending on the host cell or tissue and/or the site of integration of the sequence. In a host cell containing such a target sequence and

5 expressing an inhibitor fusion protein of the invention, transcription of the target sequence can be down regulated in a controlled manner by altering the concentration of Tc (or analogue) in contact with the host cell. For example, when the inhibitor fusion protein binds to *tetO* in the absence of Tc, the concentration of Tc in contact with the host cell is reduced to inhibit expression of the target gene. Preferably, a host cell is cultured in the absence of Tc to
10 keep target gene expression repressed. Likewise, Tc is not administered to a host organism to keep target gene expression repressed. Alternatively, when the inhibitor fusion protein binds to *tetO* in the presence of Tc, the concentration of Tc in contact with the host cell is increased to inhibit expression of the target gene. For example, Tc is added to the culture medium of a host cell or Tc is administered to a host organism to repress target gene expression.

15 The inhibitor fusion proteins described herein can inhibit a *tetO*-linked gene of interest in which the *tetO* sequences are positioned 5' of a minimal promoter sequence (e.g., tetracycline-regulated transcription units as described in Section III, above). Furthermore, the inhibitor fusion protein may be used to inhibit expression of a gene of interest in which *tetO*-linked sequences are located 3' of the promoter sequence but 5' of the transcription start site.
20 Still further, the inhibitor fusion protein may be used to inhibit expression of a gene of interest in which *tetO*-linked sequences are located 3' of the transcription start site.

Various Tc analogues as described in Section VI, part A, above, with respect to the transactivator fusion proteins can similarly be used to regulate the activity of the inhibitor fusion proteins. Moreover, the methods of *in vitro* culture with Tc (or analogue) and *in vivo*
25 administration of Tc (or analogue) described in Section VI, part A, are equally applicable to the transcriptional inhibitor fusion proteins.

C. Combined Positive and Negative Regulation of Gene Expression

In addition to regulating gene expression using either a transcriptional activator or
30 inhibitor fusion protein alone, the two types of fusion proteins can be used in combination to allow for both positive and negative regulation of expression of one or more target genes in a host cell. Thus, a transcriptional inhibitor protein that binds to *tetO* either (i) in the absence, but not the presence, of Tc, or (ii) in the presence, but not the absence, of Tc, can be used in combination with a transactivator protein that binds to *tetO* either (i) in the absence, but not
35 the presence, of Tc, or (ii) in the presence, but not the absence, of Tc. Transactivator proteins that bind to *tetO* in the absence, but not the presence, of Tc (e.g., wild-type TetR-activator fusion proteins) are described in further detail in USSN 08/076,726, USSN 08/076,327 and USSN 08/260,452. Transactivator fusion proteins that bind to *tetO* in the presence, but not the absence, of Tc (e.g., mutated TetR-activator fusion proteins) are described herein (see

Section I above) and in USSN 08/270,637 and USSN 08/275,876. Transcriptional inhibitor fusion proteins are described herein in Section IV.

As described above in Section III, Part C, when more than one TetR fusion protein is expressed in a host cell or organism, additional steps may be taken to inhibit heterodimerization between the different TetR fusion proteins. For example, a transactivator composed of a TetR of one class may be used in combination with a transcriptional inhibitor composed of a TetR of a second, different class that does not heterodimerize with the first class of TetR. Alternatively, amino acid residues of the TetR involved in dimerization may be mutated to inhibit heterodimerization. However, even if some heterodimerization between transactivator and inhibitor fusion proteins occurs in a host cell, sufficient amounts of homodimers should be produced to allow for efficient positive and negative regulation as described herein.

It will be appreciated by those skilled in the art that various combinations of activator and inhibitor proteins can be used to regulate a single *tetO*-linked gene of interest in both a positive and negative manner or to regulate multiple *tetO*-linked genes of interest in a coordinated manner or in an independent manner using the teachings described herein. The precise regulatory components utilized will depend upon the genes to be regulated and the type of regulation desired. Several non-limiting examples of how the transactivator and inhibitor fusion proteins may be used in combination are described further below. However, many other possible combinations will be evident to the skilled artisan in view of the teachings herein and are intended to be encompassed by the invention.

In a preferred embodiment, illustrated schematically in Figure 10, expression of a *tetO*-linked target gene of interest in a host cell is regulated in both a negative and positive manner by the combination of an inhibitor fusion protein that binds to *tetO* in the absence, but not the presence, of tetracycline or analogue thereof (referred to as a tetracycline controlled silencing domain, or tSD) and an activator fusion protein that binds to *tetO* in the presence, but not the absence, of tetracycline or analogue thereof (referred to as a reverse tetracycline controlled transactivator, or rTA). In addition to *tetO* sequences, the target gene is linked to a promoter, and may contain other positive regulatory elements (e.g., enhancer sequences) that contribute to basal level, constitutive transcription of the gene in the host cell. Binding of tSD to the *tetO* sequences in the absence of tetracycline or analogue (e.g., doxycycline) inhibits the basal constitutive transcription of the gene of interest, thus keeping the gene of interest in a repressed state until gene expression is desired. When gene expression is desired, the concentration of tetracycline or analogue (e.g., doxycycline) in contact with the host cell increased. Upon addition of the drug, tSD loses the ability to bind to *tetO* sequences whereas the previously unbound rTA acquires the ability to bind to *tetO* sequences. The resultant binding of rTA to the *tetO* sequences linked to the gene of interest thus stimulates transcription of the gene of interest. The level of expression may be controlled by the concentration of tetracycline or analogue, the type of Tc analogue used, the

duration of induction, etc., as described previously herein. It will be appreciated that the reverse combination of fusion proteins (i.e., the inhibitor binds in the presence but not the absence of the drug and the activator binds in the absence but not the presence of the drug) can also be used. In this case, expression of the gene of interest is kept repressed by
5 contacting the host cell with the drug (e.g., culture with Tc or analogue) and gene expression is activated by removal of the drug.

In another embodiment, the activator and inhibitor fusion proteins, as described in the previous paragraph, are used in combination to coordinately regulate, in both a positive and negative manner, two genes of interest using the bidirectional *tetO*-linked transcription unit
10 described in Section III, Part B above. In this case, Gene 1 and Gene 2 are linked to the same *tetO* sequence(s), but in opposite orientations. The inhibitor fusion protein is used to repress basal levels of transcription of both Gene 1 and Gene 2 in a coordinate manner, whereas the transactivator fusion protein is used to stimulate expression of Gene 1 and Gene 2 in a coordinate manner.

15 In yet another embodiment, the activator and inhibitor fusion proteins are used to independently regulate two or more genes of interest using the *tetO*-linked transcription units as described in Section III, Part C above. For example, in one embodiment, a transactivator fusion protein that binds to one class of *tetO* sequences (e.g., class A) in the presence, but not the absence of Tc or analogue is used in combination with an inhibitor fusion protein that
20 binds to a second, different class of *tetO* sequences (e.g., class B) also in the presence, but not the absence, of Tc or analogue. In a host cell containing Gene 1 linked to class A *tetO* sequences and Gene 2 linked to class B *tetO* sequences, both genes will be expressed at basal levels in the absence of the drug, whereas expression of Gene 1 will be stimulated upon addition of the drug and expression of Gene 2 will be repressed upon addition of the drug.

25 Alternatively, in another embodiment, the transactivator binds to one class of *tetO* sequences (e.g., class A) in the presence, but not the absence, of Tc or analogue and the inhibitor fusion protein binds to a second, different class of *tetO* sequences (e.g., class B) in the absence but not the presence of Tc or analogue. In the host cell as described in the previous paragraph, Gene 1 will be expressed at basal levels in the absence of the drug and
30 will be stimulated upon addition of the drug, whereas Gene 2 will be repressed in the absence of the drug but will have basal levels expression upon addition of the drug. Various other possible combinations will be apparent to the skilled artisan. Transactivator and inhibitor fusion proteins that bind to different classes of *tetO* sequences can be prepared as described in Section I, Part A. Target transcription units comprising *tetO* sequences of different classes
35 can be prepared as described in Section III, Part C.

VII. Applications of the Invention

The invention is widely applicable to a variety of situations where it is desirable to be able to turn gene expression on and off, or regulate the level of gene expression, in a rapid, efficient and controlled manner without causing pleiotropic effects or cytotoxicity. Thus, the system of the invention has widespread applicability to the study of cellular development and differentiation in eukaryotic cells, plants and animals. For example, expression of oncogenes can be regulated in a controlled manner in cells to study their function. Additionally, the system can be used to regulate the expression of site-specific recombinases, such as CRE or FLP, to thereby allow for irreversible modification of the genotype of a transgenic organism under controlled conditions at a particular stage of development. For example, drug resistance markers inserted into the genome of transgenic plants that allow for selection of a particular transgenic plant could be irreversibly removed via a Tc-regulated site specific recombinase. Other applications of the regulatory system of the invention include:

A. Gene Therapy

The invention may be particularly useful for gene therapy purposes, in treatments for either genetic or acquired diseases. The general approach of gene therapy involves the introduction of nucleic acid into cells such that one or more gene products encoded by the introduced genetic material are produced in the cells to restore or enhance a functional activity. For reviews on gene therapy approaches see Anderson, W.F. (1992) *Science* 256:808-813; Miller, A.D. (1992) *Nature* 357:455-460; Friedmann, T. (1989) *Science* 244:1275-1281; and Courmoyer, D., et al. (1990) *Curr. Opin. Biotech.* 1:196-208. However, current gene therapy vectors typically utilize constitutive regulatory elements which are responsive to endogenous transcription factors. These vector systems do not allow for the ability to modulate the level of gene expression in a subject. In contrast, the inducible regulatory system of the invention provides this ability.

To use the system of the invention for gene therapy purposes, in one embodiment, cells of a subject in need of gene therapy are modified to contain 1) nucleic acid encoding a transactivator fusion protein of the invention in a form suitable for expression of the transactivator in the host cells and 2) a gene of interest (e.g., for therapeutic purposes) operatively linked to a *tet* operator sequence(s). The cells of the subject can be modified *ex vivo* and then introduced into the subject or the cells can be directly modified *in vivo* (methods for modification of the cells are described above in Section II). Expression of the gene of interest in the cells of the subject is then stimulated by administering Tc or a Tc analogue to the patient. The level of gene expression can be varied depending upon which particular Tc analogue is used as the inducing agent. The level of gene expression can also be modulated by adjusting the dose of the tetracycline, or analogue thereof, administered to

the patient to thereby adjust the concentration achieved in the circulation and the tissues of interest.

Moreover, in another embodiment, a transcriptional inhibitor fusion protein is used to further control the level of expression of the gene of interest. For example, the cells of the subject can be modified to also contain a nucleic acid encoding a transcriptional inhibitor fusion protein that binds to *tetO* in the absence of Tc. The nucleic acid is in a form suitable for expression of the inhibitor fusion protein in the host cells. Thus, prior to administration of Tc (or analogue) to the subject, the basal level of transcription of the gene of interest will be kept silent by the inhibitor fusion protein. Upon administration of Tc, binding of the inhibitor fusion protein to *tetO* will be inhibited whereas binding of the transactivator fusion will be induced, thereby stimulating transcription of the gene of interest. Such combined positive and negative regulation of gene expression using both a transactivator fusion protein and transcriptional inhibitor fusion protein of the invention is illustrated schematically in Figure 10.

Conventional detection methods known in the art, such as an enzyme linked immunosorbent assay, can be used to monitor the expression of the regulated protein of interest in the host cells and the concentration of Tc or Tc analogue can be varied until the desired level of expression of the protein of interest is achieved. Accordingly, expression of a protein of interest can be adjusted according to the medical needs of an individual, which may vary throughout the lifetime of the individual. To stop expression of the gene of interest in cells of the subject, administration of the inducing agent is stopped. Thus, the regulatory system of the invention offers the advantage over constitutive regulatory systems of allowing for modulation of the level of gene expression depending upon the requirements of the therapeutic situation.

Genes of particular interest to be expressed in cells of a subject for treatment of genetic or acquired diseases include those encoding adenosine deaminase, Factor VIII, Factor IX, dystrophin, β -globin, LDL receptor, CFTR, insulin, erythropoietin, anti-angiogenesis factors, growth hormone, glucocerebrosidase, β -glucuronidase, α 1-antitrypsin, phenylalanine hydroxylase, tyrosine hydroxylase, ornithine transcarbamylase, arginosuccinate synthetase, UDP-glucuronosyl transferase, apoA1, TNF, soluble TNF receptor, interleukins (e.g., IL-2), interferons (e.g., α - or γ -IFN) and other cytokines and growth factors. Cells types which can be modified for gene therapy purposes include hematopoietic stem cells, myoblasts, hepatocytes, lymphocytes, skin epithelium and airway epithelium. For further descriptions of cell types, genes and methods for gene therapy see e.g., Wilson, J.M et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:3014-3018; Armentano, D. et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6141-6145; Wolff, J.A. et al. (1990) *Science* 247:1465-1468; Chowdhury, J.R. et al. (1991) *Science* 254:1802-1805; Ferry, N. et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8377-8381; Wilson, J.M. et al. (1992) *J. Biol. Chem.* 267:963-967; Quantin, B. et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:2581-2584; Dai, Y. et al. (1992)

- Proc. Natl. Acad. Sci. USA* 89:10892-10895; van Beusechem, V.W. et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:7640-7644; Rosenfeld, M.A. et al. (1992) *Cell* 68:143-155; Kay, M.A. et al. (1992) *Human Gene Therapy* 3:641-647; Cristiano, R.J. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:2122-2126; Hwu, P. et al. (1993) *J. Immunol.* 150:4104-4115; and Herz, J. and Gerard, R.D. (1993) *Proc. Natl. Acad. Sci. USA* 90:2812-2816.

Gene therapy applications of particular interest in cancer treatment include overexpression of a cytokine gene (e.g., TNF- α) in tumor infiltrating lymphocytes or ectopic expression of cytokines in tumor cells to induce an anti-tumor immune response at the tumor site), expression of an enzyme in tumor cells which can convert a non-toxic agent into a toxic agent, expression of tumor specific antigens to induce an anti-tumor immune response, expression of tumor suppressor genes (e.g., p53 or Rb) in tumor cells, expression of a multidrug resistance gene (e.g., MDR1 and/or MRP) in bone marrow cells to protect them from the toxicity of chemotherapy.

Gene therapy applications of particular interest in treatment of viral diseases include expression of trans-dominant negative viral transactivation proteins, such as trans-dominant negative tat and rev mutants for HIV or trans-dominant ICp4 mutants for HSV (see e.g., Balboni, P.G. et al. (1993) *J. Med. Virol.* 41:289-295; Liem, S.E. et al. (1993) *Hum. Gene Ther.* 4:625-634; Malim, M.H. et al. (1992) *J. Exp. Med.* 176:1197-1201; Daly, T.J. et al. (1993) *Biochemistry* 32:8945-8954; and Smith, C.A. et al. (1992) *Virology* 191:581-588), expression of trans-dominant negative envelope proteins, such as env mutants for HIV (see e.g., Steffy, K.R. et al. (1993) *J. Virol.* 67:1854-1859), intracellular expression of antibodies, or fragments thereof, directed to viral products ("internal immunization", see e.g., Marasco, W.A. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:7889-7893) and expression of soluble viral receptors, such as soluble CD4. Additionally, the system of the invention can be used to conditionally express a suicide gene in cells, thereby allowing for elimination of the cells after they have served an intended function. For example, cells used for vaccination can be eliminated in a subject after an immune response has been generated the subject by inducing expression of a suicide gene in the cells by administering Tc or a Tc analogue to the subject.

The Tc-controlled regulatory system of the invention has numerous advantages properties that it particularly suitable for application to gene therapy. For example, the system provides an "on"/"off" switch for gene expression that allows for regulated dosaging a gene product in a subject. There are several situations in which it may be desirable to be able to provide a gene product at specific levels and/or times in a regulated manner, rather than simply expressing the gene product constitutively at a set level. For example, a gene of interest can be switched "on" at fixed intervals (e.g., daily, alternate days, weekly, etc.) to provide the most effective level of a gene product of interest at the most effective time. The level of gene product produced in a subject can be monitored by standard methods (e.g., direct monitoring using an immunological assay such as ELISA or RIA or indirectly by monitoring of a laboratory parameter dependent upon the function of the gene product of

interest, e.g., blood glucose levels and the like). This ability to turn "on" expression of a gene at discrete time intervals in a subject while also allowing for the gene to be kept "off" at other times avoids the need for continued administration of a gene product of interest at intermittent intervals. This approach avoids the need for repeated injections of a gene product, which may be painful and/or cause side effects and would likely require continuous visits to a physician. In contrast, the system of the invention avoids these drawbacks. Moreover, the ability to turn "on" expression of a gene at discrete time intervals in a subject allows for focused treatment of diseases which involve "flare ups" of activity (e.g., many autoimmune diseases) only at times when treatment is necessary during the acute phase when pain and symptoms are evident. At times when such diseases are in remission, the expression system can be kept in the "off" state.

Gene therapy applications that may particularly benefit from this ability to modulate gene expression during discrete time intervals include the following non-limiting examples:

Rheumatoid arthritis - genes which encode gene products that inhibit the production of inflammatory cytokines (e.g., TNF, IL-1 and IL-12). can be expressed in subjects. Examples of such inhibitors include soluble forms of a receptor for the cytokine. Additionally or alternatively, the cytokines IL-10 and/or IL-4 (which stimulate a protective Th2-type response) can be expressed. Moreover, a glucocorticomimetic receptor (GCMR) can be expressed.

Hypopituitarism - the gene for human growth hormone can be expressed in such subjects only in early childhood, when gene expression is necessary, until normal stature is achieved, at which time gene expression can be downregulated.

Wound healing/Tissue regeneration - Factors (e.g., growth factors, angiogenic factors, etc.) necessary for the healing process can be expressed only when needed and then downregulated.

Anti-Cancer Treatments - Expression of gene products useful in anti-cancer treatment can be limited to a therapeutic phase until retardation of tumor growth is achieved, at which time expression of the gene product can be downregulated. Possible systemic anti-cancer treatments include use of tumor infiltrating lymphocytes which express immunostimulatory molecules (e.g., IL-2, IL-12 and the like), angiogenesis inhibitors (PF4, IL-12, etc.), Her-regulin, Leukoregulin (see PCT Publication No. WO 85/04662), and growth factors for bone marrow support therapy, such as G-CSF, GM-CSF and M-CSF. Regarding the latter, use of the regulatory system of the invention to express factors for bone marrow support therapy allows for simplified therapeutic switching at regular intervals from chemotherapy to bone marrow support therapy (similarly, such an approach can also be applied to AIDS treatment, e.g., simplified switching from anti-viral treatments to bone marrow support treatment). Furthermore, controlled local targeting of anti-cancer treatments are also possible. For example, expression of a suicide gene by a regulator of the invention, wherein the regulator

itself is controlled by, for example, a tumor-specific promoter or a radiation-induced promoter.

5 In another embodiment, the regulatory system of the invention is used to express angiogenesis inhibitor(s) from within a tumor via a transgene regulated by the system of the invention. Expression of angiogenesis inhibitors in this manner may be more efficient than systemic administration of the inhibitor and would avoid any deleterious side effects that might accompany systemic administration. In particular, restricting angiogenesis inhibitor expression to within tumors could be particularly useful in treating cancer in children still undergoing angiogenesis associated with normal cell growth.

10 In another embodiment, high level regulated expression of cytokines may represent a method for focusing a patient's own immune response on tumor cells. Tumor cells can be transduced to express chemoattractant and growth promoting cytokines important in increasing an individual's natural immune response. Because the highest concentrations of cytokines will be in the proximity of the tumor, the likelihood of eliciting an immunological response to tumor antigens is increased. A potential problem with this type of therapy is that those tumor cells producing the cytokines will also be targets of the immune response and therefor the source of the cytokines will be eliminated before eradication of all tumor cells can be certain. To combat this, expression of viral proteins known to mask infected cells from the immune system can be placed under regulation, along with the cytokine gene(s), in 20 the same cells. One such protein is the E19 protein from adenovirus (see e.g., Cox, *Science* 247:715). This protein prevents transport of class I HLA antigens to the surface of the cell and hence prevents recognition and lysis of the cell by the host's cytotoxic T cells. Accordingly, regulated expression of E19 in tumor cells could shield cytokine producer cells from cytotoxic T cells during the onset of an immune response provoked by cytokine 25 expression. After a sufficient period of time has elapsed to eradicate all tumor cells but those expressing E19, E19 expression can be turned off, causing these cells then to fall victim to the provoked anti-tumor immune response.

30 Benign prostatic hypertrophy - Similar to the above, a suicide gene can be regulated by a regulator of the invention, wherein the regulator itself is controlled by, for example, a prostate-specific promoter.

The ability to express a suicide gene (e.g., an apoptosis gene, TK gene, etc) in a controlled manner using the regulatory system of the invention adds to the general safety and usefulness of the system. For example, at the end of a desired therapy, expression of a suicide gene can be triggered to eliminate cells carrying the gene therapy vector, such as cells 35 in a bioinert implant, cells that have disseminated beyond the intended original location, etc. Moreover, if a transplant becomes tumorous or has side effects, the cells can be rapidly eliminated by induction of the suicide gene. The use of more than one Tc-controlled "on"/"off" switch in one cell allows for completely independent regulation of a suicide gene compared to regulation of a gene of therapeutic interest (as described in detail herein).

The regulatory system of the invention further offers the ability to establish a therapeutically relevant expression level for a gene product of interest in a subject, in contrast to unregulated constitutive expression which offers no flexibility in the level of gene product expression that can be achieved. A physiologically relevant level of gene product expression can be established based on the particular medical need of the subject, e.g., based on laboratory tests that monitor relevant gene product levels (using methods as described above). In addition to the clinical examples and gene products already discussed above with gene to dosaging of the gene product, other therapeutically relevant gene products which can be expressed at a desired level at a desired time include: Factor XIII and IX in hemophiliacs (e.g., expression can be elevated during times of risk of injury, such as during sports); insulin or amylin in diabetics (as needed, depending on the state of disease in the subject, diet, etc.); erythropoietin to treat erythrocytopenia (as needed, e.g., at end-stage renal failure); low-density lipoprotein receptor (LDLr) or very low-density lipoprotein receptor (VLDLr) for arterosclerosis or gene therapy in liver (e.g., using *ex vivo* implants). Applications to treatment of central nervous system disorders are also encompassed. For example, in Alzheimer's disease, "fine tuned" expression of choline acetyl transferase (ChAT) to restore acetylcholine levels, neurotrophic factors (e.g., NGF, BDNF and the like) and/or complement inhibitors (e.g., sCR1, sMCP, sDAF, sCD59 etc.) can be accomplished. Such gene products can be provided, for example, by transplanted cells expressing the gene products in a regulated manner using the system of the invention. Moreover, Parkinson's disease can be treated by "fine tuned" expression of tyrosine hydroxylase (TH) to increase levodopa and dopamine levels.

In addition to the proteinaceous gene products discussed above, gene products that are functional RNA molecules (such as anti-sense RNAs and ribozymes) can be expressed in a controlled manner in a subject for therapeutic purposes. For example, a ribozyme can be designed which discriminates between a mutated form of a gene and a wild-type gene. Accordingly, a "correct" gene (e.g., a wild-type p53 gene) can be introduced into a cell in parallel with introduction of a regulated ribozyme specific for the mutated form of the gene (e.g., a mutated endogenous p53 gene) to remove the defective mRNA expressed from the endogenous gene. This approach is particularly advantageous in situations in which a gene product from the defective gene would interfere with the action of the exogenous wild-type gene.

Expression of a gene product in a subject using the regulatory system of the invention is modulated using tetracycline or analogues thereof. Such drugs can be administered by any route appropriate for delivery of the drug to its desired site of action (e.g., delivery to cells containing a gene whose expression is to be regulated). Depending on the particular cell types involved, preferred routes of administration may include oral administration, intravenous administration and topical administration (e.g., using a transdermal patch to reach

cells of a localized transplant under the skin, such as keratinocytes, while avoiding any possible side effects from systemic treatment).

In certain gene therapy situations, it may be necessary or desirable to take steps to avoid or inhibit unwanted immune reactions in a subject receiving treatment. To avoid a reaction against the cells expressing the therapeutic gene product, a subject's own cells are generally used, when possible, to express the therapeutic gene product, either by *in vivo* modification of the subject's cells or by obtaining cells from the subject, modifying them *ex vivo* and returning them to the subject. In situations where allogeneic or xenogeneic cells are used to express a gene product of interest, the regulatory system of the invention, in addition to regulating a therapeutic gene, can also be used to regulate one or more genes involved in the immune recognition of the cells to inhibit an immune reaction against the foreign cells. For example, cell-surface molecules involved in recognition of a foreign cell by T lymphocytes can be downmodulated on the surface of a foreign cell used for delivery of a therapeutic gene product, such as by regulated expression in the foreign cell of a ribozyme which cleaves the mRNA encoding the cell-surface molecule. Particularly preferred cell surface molecules which can be downmodulated in this manner to inhibit an unwanted immune response include class I and/or class II major histocompatibility complex (MHC) molecules, costimulatory molecules (e.g., B7-1 and/or B7-2), CD40, and various "adhesion" molecules, such as ICAM-1 or ICAM-2. Using approaches described herein for independent but coordinate regulation of multiple genes in the same cell, the down-regulation of expression of a cell-surface molecule(s) in a host cell can be coordinated with the up-regulation of expression of a therapeutic gene. Accordingly, after therapy is completed and expression of the therapeutic gene is halted, expression of the endogenous cell surface molecule(s) can be restored to normal.

Furthermore, as described above regarding anti-cancer treatments, a viral protein (e.g., adenovirus E19 protein) that downmodulates expression of MHC antigens can be regulated in host cells using the system of the invention as a means of avoiding unwanted immunological reactions.

In addition to avoiding or inhibiting an immune response against a foreign cell delivering a therapeutic gene product, it may also be necessary, in certain situations, to avoid or inhibit an immune response against certain components of the regulatory system of the invention (e.g., the regulator fusion proteins described herein) that are expressed in a subject, since these fusion proteins contain non-mammalian polypeptides that may stimulate an unwanted immune reaction. In this regard, regulator fusion proteins can be designed and/or selected for a decreased ability to stimulate an immune response in a host. For example, a transcriptional activator domain for use in the regulator fusion protein can be chosen which has minimal immunogenicity. In this regard, a wild-type transcriptional activation domain of the herpes simplex virus protein VP16 may not be a preferred transcriptional activation domain for use *in vivo*, since it may stimulate an immune response in mammals. Alternative

transcriptional activation domains can be used, as described herein, based on their reduced immunogenicity in a subject. For example, a transcriptional activation domain of a protein of the same species as the host may be preferred (e.g., a transcriptional activation domain from a human protein for use of a regulatory fusion protein in humans). Alternatively, a regulatory fusion protein of the invention can be modified to reduce its immunogenicity in subjects, e.g., by identifying and modifying one or more dominant T cell epitopes within a polypeptide of the fusion protein (e.g., either the Tet repressor moiety or the transcriptional modulator moiety, such as a VP16 polypeptide). Such T cell epitopes can be identified by standard methods and altered by mutagenesis, again by standard methods. A modified form of a regulator fusion protein can then be selected which retains its original transcriptional regulatory ability yet which exhibits reduced immunogenicity in a subject as compared to an unmodified fusion protein.

In addition to the foregoing, all conventional methods for generally or specifically downmodulating immune responses in subjects can be combined with the use of the regulatory system of the invention in situations where inhibition of immune responses is desired. General immunosuppressive agents, such as cyclosporin A and/or FK506, can be administered to the subject. Alternatively, immunomodulatory agents which may allow for more specific immunosuppression can be used. Such agents may include inhibitors of costimulatory molecules (e.g., a CTLA4Ig fusion protein, soluble CD4, anti-CD4 antibodies, anti-B7-1 and/or anti-B7-2 antibodies or anti-gp39 antibodies).

Finally, in certain situations, a delivery vehicle for cells expressing a therapeutic gene can be chosen which minimizes exposure of transplanted cells to the immune system. For example, cells can be implanted into bioinert capsules/biocompatible membranes with pores which allow for diffusion of proteins (e.g., a therapeutic gene product of interest) out of the implant and diffusion of nutrients and oxygen into the implant but which prevent entry of immune cells, thereby avoiding exposure of the transplanted cells to the immune system (as has been applied to islet cell transplantation).

B. Production of Proteins in Vitro

Large scale production of a protein of interest can be accomplished using cultured cells *in vitro* which have been modified to contain 1) a nucleic acid encoding a transactivator fusion protein of the invention in a form suitable for expression of the transactivator in the cells and 2) a gene encoding the protein of interest operatively linked to a *tet* operator sequence(s). For example, mammalian, yeast or fungal cells can be modified to contain these nucleic acid components as described herein. The modified mammalian, yeast or fungal cells can then be cultured by standard fermentation techniques in the presence of Tc or an analogue thereof to induce expression of the gene and produce the protein of interest. Accordingly, the invention provides a production process for isolating a protein of interest. In the process, a host cell (e.g., a yeast or fungus), into which has been introduced both a nucleic acid

encoding a transactivator fusion protein of the invention and a nucleic acid encoding the protein of the interest operatively linked to at least one *tet* operator sequence, is grown at production scale in a culture medium in the presence of tetracycline or a tetracycline analogue to stimulate transcription of the nucleotides sequence encoding the protein of interest (i.e., the nucleotide sequence operatively linked to the *tet* operator sequence(s)) and the protein of interest is isolated from harvested host cells or from the culture medium. Standard protein purification techniques can be used to isolate the protein of interest from the medium or from the harvested cells.

10 C. Production of Proteins in Vivo

The invention also provides for large scale production of a protein of interest in animals, such as in transgenic farm animals. Advances in transgenic technology have made it possible to produce transgenic livestock, such as cattle, goats, pigs and sheep (reviewed in Wall, R.J. et al. (1992) *J. Cell. Biochem.* 49:113-120; and Clark, A.J. et al. (1987) *Trends in Biotechnology* 5:20-24). Accordingly, transgenic livestock carrying in their genome the components of the inducible regulatory system of the invention can be constructed, wherein a gene encoding a protein of interest is operatively linked to at least one *tet* operator sequence. Gene expression, and thus protein production, is induced by administering Tc (or analogue thereof) to the transgenic animal. Protein production can be targeted to a particular tissue by linking the nucleic acid encoding the transactivator fusion protein to an appropriate tissue-specific regulatory element(s) which limits expression of the transactivator to certain cells. For example, a mammary gland-specific regulatory element, such as the milk whey promoter (U.S. Patent No. 4,873,316 and European Application Publication No. 264,166), can be linked to the transactivator transgene to limit expression of the transactivator to mammary tissue. Thus, in the presence of Tc (or analogue), the protein of interest will be produced in the mammary tissue of the transgenic animal. The protein can be designed to be secreted into the milk of the transgenic animal, and if desired, the protein can then be isolated from the milk.

30 D. Animal Models of Human Disease

The transcriptional activator and inhibitor proteins of the invention can be used alone or in combination to stimulate or inhibit expression of specific genes in animals to mimic the pathophysiology of human disease to thereby create animal models of human disease. For example, in a host animal, a gene of interest thought to be involved in a disease can be placed under the transcriptional control of one or more *tet* operator sequences (e.g., by homologous recombination, as described herein). Such an animal can be mated to a second animal carrying one or more transgenes for a transactivator fusion protein and/or an inhibitor fusion protein to create progeny that carry both a tetracycline-regulated fusion protein(s) gene and a tet-regulated target sequence. Expression of the gene of interest in these progeny can be

modulated using tetracycline (or analogue). For example, expression of the gene of interest can be downmodulated using a transcriptional inhibitor fusion protein to examine the relationship between gene expression and the disease. Such an approach may be advantageous over gene "knock out" by homologous recombination to create animal models of disease, since the tet-regulated system described herein allows for control over both the levels of expression of the gene of interest and the timing of when gene expression is down- or up-regulated.

E. Production of Stable Cell Lines for Gene Cloning and Other Uses

The transcriptional inhibitor system described herein can be used keep gene expression "off" (i.e., expressed) to thereby allow production of stable cell lines that otherwise may not be produced. For example, stable cell lines carrying genes that are cytotoxic to the cells can be difficult or impossible to create due to "leakiness" in the expression of the toxic genes. By repressing gene expression of such toxic genes using the transcriptional inhibitor fusion proteins of the invention, stable cell lines carrying toxic genes may be created. Such stable cell lines can then be used to clone such toxic genes (e.g., inducing the expression of the toxic genes under controlled conditions using Tc or analog). General methods for expression cloning of genes, to which the transcriptional inhibitor system of the invention can be applied, are known in the art (see e.g., Edwards, C. P. and Aruffo, A. (1993) *Curr. Opin. Biotech.* 4:558-563). Moreover, the transcriptional inhibitor system can be applied to inhibit basal expression of genes in other cells to create stable cell lines, such as in embryonic stem (ES) cells. Residual expression of certain genes introduced into ES stems may result in an inability to isolate stably transfected clones. Inhibition of transcription of such genes using the transcriptional inhibitor system described herein may be useful in overcoming this problem.

Advantages

The inducible regulatory system of the invention utilizing a transactivator fusion protein addresses and overcomes many of the limitations of other inducible regulatory systems in the art. For example, very high intracellular concentrations of the transcriptional activator fusion protein of the invention are not required for efficient regulation of gene expression. Additionally, since gene expression is induced by adding rather than removing the inducing agent, the induction kinetics in the system of the invention are not limited by the rate of removal of the inducing agent and thus are typically faster. Moreover, the inducing agent is only present when gene transcription is induced, thereby avoiding the need for the continuous presence of an agent to keep gene expression off.

Use of the transcriptional inhibitor fusion proteins of the invention to inhibit transcription in eukaryotic cells also provide advantages over the use of prokaryotic repressors alone (e.g., TetR, lacR) to inhibit transcription in eukaryotic cells. Since the

inhibitor fusion proteins of the invention contain a eukaryotic transcriptional silencer domain, these fusion proteins should be more efficient at repressing transcription in eukaryotic cells, and thus may potentially require lower intracellular concentrations for efficient repression with less likelihood of "leakiness". Additionally, by insertion of *tetO* sequences into the regulatory region of an endogenous gene, the transcriptional inhibitor fusion proteins of the invention can be used to down-regulate constitutive and/or tissue-specific expression of endogenous genes.

Furthermore, in contrast to various versions of the *lac* system (e.g., Labow et al. (1990) *Mol. Cell. Biol.* 10:3343-3356; Baim et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:5072-5076), which are limited by the negative properties of the inducing agent (IPTG) and/or by the need to increase the temperature in order to induce gene expression (which may elicit pleiotropic effects), the inducing agent used in the system of the invention (Tc or an analogue thereof) has many advantageous properties: 1) Tc and analogues thereof exhibit high affinity for TetR and low toxicity for eukaryotic cells, and thus can be used for gene induction at concentrations that do not affect cell growth or morphology; 2) Tc analogues which retain TetR binding but which have reduced antibiotic activity exist and can be used as inducing agents, thereby avoiding possible side effects from the antibiotic property of Tc; 3) the pharmacokinetic properties of Tc and Tc analogues enable rapid and efficient cellular uptake and penetration of physiological barriers, such as the placenta or the blood-brain barrier; and 4) Tc analogues with different induction capabilities permit modulation of the level of gene expression.

Thus, the invention provides an inducible regulatory system which allows for rapid activation of gene transcription without cellular toxicity and a range of induction indices. The increase in gene expression upon induction typically is between 1000- and 2000-fold and can be as high as about 20,000-fold. Alternatively, lower levels of gene induction, e.g., 10-fold, can be achieved depending upon which inducing agent is used. This system can be utilized in a wide range of applications. These applications include gene therapy, large-scale production of proteins in cultured cells or in transgenic farm animals, and the study of gene function, for example in relationship to cellular development and differentiation. Moreover, the novel transcription units of the invention allow for coordinate or independent regulation of the expression of multiple genes utilizing the regulatory components of the invention.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

EXAMPLE 1: Selection of a Mutated Tet Repressor and Construction of a Tetracycline Inducible Transcriptional Activator

A "reverse" Tet repressor, which binds to its target DNA in the presence rather than the absence of tetracycline, was generated by chemical mutagenesis and selection essentially as described in Hecht, B. et al. (1993) *J. Bacteriology* 175:1206-1210. Single-stranded DNA (coding and non-coding strands) encoding the wild-type Tn10-derived Tet repressor was chemically mutagenized with sodium nitrite. Single-stranded DNAs (40 µg in 40 µl in Tris-EDTA buffer) were mixed with 10 µl of 2.5 M sodium acetate (pH 4.3) and 50 µl of sodium nitrate ranging between 0.25 M and 2 M and incubated for 45 to 60 minutes at room temperature. After mutagenesis, the complementary strand was synthesized using reverse transcriptase or by amplification using the polymerase chain reaction with Taq DNA polymerase. Since the mutagenesis procedure yields multiple mutations in the DNA, three fragments of the gene, of about 200 base pairs each, were individually subcloned into a wt Tet repressor gene in a recombinant expression vector to replace the corresponding portion of the wild-type gene. This created a pool of mutated Tet repressor genes wherein each gene had mostly single mutations in the 200 base pair mutagenized fragment of the gene.

The pool of mutated Tet repressors were screened in a genetic assay which positively selects for a functional interaction between a Tet repressor and its cognate operator using *E. coli* strain WH207(λWH25) (the construction of this strain is described in detail in Wissmann, A. et al. (1991) *Genetics* 128:225-232). In this *E. coli* strain, *tet* operators direct the expression of divergently arranged β-galactoside (*lacZ*) and Lac repressor (*lacI*) genes and the *lac* regulatory region directs the expression of a galactokinase (*galK*) gene. Binding of Tet repressors to *tet* operators turns off transcription of the *lacI* and *lacZ* genes. The absence of Lac repressor allows for expression of the *galK* gene, which enables the *E. coli* strain to use galactose as a sole carbon source, which serves as one marker. The *lacZ*⁻ phenotype serves as a second marker. Thus, bacteria containing Tet repressors which bind to *tet* operators have a Gal⁺, *lacZ*⁻ phenotype. Bacteria containing wild-type Tet repressors have a Gal⁺, *lacZ*⁻ phenotype in the absence of tetracycline. A mutated "reverse" Tet repressor (rTetR) was selected based upon a Gal⁺, *lacZ*⁻ phenotype in the presence of tetracycline.

The nucleotide and amino acid sequence of the rTetR mutant are shown in SEQ ID NOs: 1 (nucleotide positions 1-621) and 2 (amino acid positions 1-207), respectively. Sequence analysis of the rTetR mutant showed the following amino acid and nucleotide changes:

| <u>aa (position)</u> | | <u>affected codon</u> | |
|----------------------|---------------|-----------------------|---------------|
| <u>wild-type</u> | <u>mutant</u> | <u>wild-type</u> | <u>mutant</u> |
| glu (71) | lys | GAA | AAA |
| asp (95) | asn | GAT | AAT |
| leu (101) | ser | TTA | TCA |
| gly (102) | asp | GGT | GAT |

Two additional mutations did not result in an amino acid exchange:

5

| <u>aa (position)</u> | | <u>affected codon</u> | |
|----------------------|---------------|-----------------------|---------------|
| <u>wild-type</u> | <u>mutant</u> | <u>wild-type</u> | <u>mutant</u> |
| leu (41) | leu | TTG | CTG |
| arg (80) | arg | CGT | CGC |

To convert the rTetR mutant to a transcriptional activator, a 399 base pair XbaI/Eco47III fragment encoding amino acids 3 to 135 of rTetR (i.e., encompassing the mutated region) was exchanged for the corresponding restriction fragment of the expression vector pUHD15-1 to create pUHD17-1. In pUHD15-1, nucleotide sequences encoding wild-type TetR are linked in frame to nucleotide sequences encoding the C-terminal 130 amino acids of herpes simplex virus VP16. These transactivator sequences are flanked upstream by a CMV promoter/enhancer and downstream by an SV40 poly(A) site (the construction of pUHD15-1 is described in more detail in USSN 08/076,726 and Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* **89**:5547-5551). Thus, in pUHD17-1, nucleotide sequences encoding the reverse TetR mutant are linked in frame to VP16 sequences to create a reverse Tc-controlled transactivator (referred to herein as tTAR^R). The analogous exchange of the mutated region of rTetR for the wild-type region of TetR was performed with plasmid pUHD152-1, which is the same as pUHD15-1 except that it additionally contains nucleotide sequences encoding a nuclear localization signal linked in-frame to the 5' end of the nucleotide sequences encoding the Tet repressor. The amino acid sequence of the nuclear localization signal is MPKRPRP (SEQ ID NO: 5), which is linked to the serine at amino acid position 2 of TetR. The resulting expression vector encoding the reverse Tc-controlled transactivator including a nuclear localization signal (referred to herein as ntTAR^R) was named pUHD172-1.

EXAMPLE 2: Tetracycline-Induced Stimulation of Transcription by tTAR^R**Transient Transfection**

The pUHD17-1 and pUHD172-1 expression vectors were transiently transfected by a standard calcium phosphate method into HeLa cells together with a reporter plasmid, pUHC13-3, in which heptameric *tet* operators are fused upstream of a minimal hCMV-promoter and a luciferase reporter gene (the reporter plasmid is described in detail in USSN 08/076,726 and Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551). After incubation of the transfected cells at 37 °C for 20 hours in the presence or absence of tetracycline (or an analogue thereof), luciferase activity was assayed as follows: Cells grown to ~80 % confluency in 35 mm dishes in Eagle's minimal essential medium were washed with 2 ml of phosphate-buffered saline before they were lysed in 25 mM Tris phosphate, pH 7.8/2 mM dithiothreitol/2 mM diaminocyclohexanetetraacetic acid/10 % glycerol/1 % Triton-X-100 for 10 minutes at room temperature. The lysate was scraped off the culture dishes and centrifuged for 10 seconds in an Eppendorf centrifuge. Next, aliquots (10 µl) of the supernatant were mixed with 250 µl of 25 mM glycylglycine/15 mM MgSO₄/5 mM ATP and assayed for luciferase activity in a Lumat LB9501 (Berthold, Wildbad, F.R.G.) using the integral mode (10 seconds). D-Luciferin (L6882, Sigma) was used at 0.5 mM. The background signal measured in extracts of HeLa cells that did not contain a luciferase gene was indistinguishable from the instrument background (80-120 relative light units (rlu)/10 sec.). Protein content of the lysate was determined according to Bradford (Bradford, M.M. (1976) *Anal. Biochem.* 72:248-254). Cells transfected with plasmids encoding either tTAR^R or ntTAR^R showed an increased level of luciferase activity in the presence of tetracyclines. This effect was consistently more pronounced when anhydrotetracycline (ATc) was used instead of tetracycline.

Stable Transfection

After this transient transfection analysis, expression vectors were prepared for stable transfection of cells. A pSV2neo-derived neomycin resistance cassette (described in Southern, P.J. and Berg, P. (1982) *J. Mol. Appl. Genet.* 1:327-341) was integrated into the transactivator expression vectors pUHD17-1 and pUHD172-1, resulting in pUHD17-1neo and pUHD172-1neo, respectively. pUHD172-1neo, coding for ntTAR^R, was stably integrated into HeLa cells by standard techniques. Ten G418-resistant cell clones were analyzed for their phenotype by transient supertransfection with pUHC13-3 carrying the luciferase gene under the control of a minimal CMV promoter and *tet* operators. Three clones, HR4, HR5 and HR10, showed a strong increase of luciferase activity in the presence of ATc. From these clones, HR5 was selected for further experiments.

To create stable transfectants for both ntTAR^R and a *tet* operator-linked luciferase reporter gene, HR5 cells were cotransfected with pUHC13-3 and pHMR272, which encodes

for hygromycin resistance (see Bernhard, H-U. et al. (1985) *Exp. Cell Res.* 158:237-243), and hygromycin resistant clones were selected. In an analogous experiment, HR5 cells were cotransfected with pUH13-7 and pHMR272. pUH13-7 contains a minimal promoter sequence spanning position +19 to -37 of the HSVtk promoter adjacent to the heptameric *tetO* sequences, rather than a minimal CMV promoter. From 21 hygromycin resistant clones, 10 showed inducible luciferase activity upon addition of Tc or doxycycline (Dc) to the culture medium. Clones containing the luciferase reporter gene linked to a minimal CMV promoter are referred to as HR5-C, whereas those containing the luciferase reporter gene linked to a minimal tk promoter are referred to HR5-T.

Six of the HR5 clones stably transfected with a ntTAR^R-dependent reporter unit and previously shown to be responsive to tetracyclines were grown in parallel in the absence or presence of 1 µg/ml doxycycline. About 3 x 10⁴ cells were plated in each 35 mm dish (4 dishes for each clone). After growth for 60 hours, cells were harvested and the luciferase activity of the extracts (in relative light units (rlu)/µg extracted protein) was determined. As shown in Table 1, the absolute expression levels of six clones demonstrate that activation of luciferase gene expression over 3 orders of magnitude is achieved in several of the double stable cell lines containing the ntTAR^R regulatory system.

It should be noted that even higher induction factors (e.g., as high as a 20,000-fold increase in expression) could be achieved if, instead of simply adding the inducing agent to the culture medium, the cells were washed prior to induction and then replated in fresh culture medium containing the inducing agent.

TABLE 1. Doxycycline-dependent luciferase activity of double stable luc+/HR5 cell clones

| Clone | Luciferase Activity, rlu/µg protein | | Induction Factor |
|---------|-------------------------------------|--------------|------------------|
| | -Doxycycline | +Doxycycline | |
| HR5-C6 | 65 | 54,911 | 845 |
| | 62 | 69,525 | 1120 |
| HR5-C11 | 100 | 165,671 | 1660 |
| | 142 | 179,651 | 1270 |
| HR5-C14 | 43 | 44,493 | 1030 |
| | 43 | 56,274 | 1310 |
| HR5-T2 | 56 | 16,696 | 298 |
| | 40 | 16,416 | 410 |
| HR5-T15 | 6.8 | 1838 | 270 |
| | 6.5 | 1688 | 260 |
| HR5-T19 | 4.8 | 1135 | 236 |
| | 5.4 | 1285 | 237 |

Induction of luciferase activity by different tetracyclines

The ability of tetracycline and several different tetracycline analogues to induce luciferase expression in HR5-C11 cells was examined. HR5-C11 cells plated at a density of about 3×10^4 cells/35 mm dish (~80% confluency). After full attachment of the cells, the following tetracyclines were added to the cultures at a concentration of 1 $\mu\text{g/ml}$: tetracycline-HCl (Tc), oxytetracycline-HCl (OTc), chlorotetracycline (CTc), anhydrotetracycline-HCl (ATc) and doxycycline-HCl (Doxy). These compounds are commercially available from Sigma Chemical Co., St. Louis, MO, and were kept in aqueous solution at a concentration of 1 $\mu\text{g/ml}$. Cells grown in the absence of antibiotic (-) served as a control. After 3 days, the cells were harvested and the luciferase activity and the protein content of the extracts were determined. The results are shown in the bar graph of Figure 1. Each bar in the figure (closed and hatched) represents the relative luciferase activity (normalized toward the amount of extracted protein) of a single culture dish. The mean of the luciferase activities obtained from the two plates grown without tetracyclines was defined as 1. Tc, CTc and OTc showed modest stimulation of luciferase activity. By contrast, ATc and Doxy stimulated luciferase activity approximately 1000 and 1500 fold, respectively.

Dose-response of luciferase activity to doxycycline in HR5-C11 cells

The above-described experiment examining the induction ability of different tetracyclines revealed that doxycycline was the most potent effector of the tetracyclines examined. Doxycycline was therefore selected to quantitatively analyze its dose-response. HR5-C11 cells were incubated with different concentrations of doxycycline and luciferase activity was measured. The data of three independent experiments are shown in Figure 2. At less than 10 ng/ml in the culture medium, doxycycline is ineffective at inducing luciferase activity. However, when the concentration was raised above 10 ng/ml, an almost linear increase in expression of luciferase was observed. Maximal activation was achieved at 1 $\mu\text{g/ml}$. At concentrations above 3 $\mu\text{g/ml}$, doxycycline showed a slight growth-inhibitory effect on HeLa cells as determined in a MTT-assay.

Kinetics of induction of the ntTA^R system

To examine the kinetics of doxycycline-induced ntTA^R-mediated induction of gene expression, the time course of induction of luciferase activity in HR5-C11 cells was monitored after addition of doxycycline to the medium (final concentration 1 $\mu\text{g/ml}$). Cells were cultured in the presence of doxycycline and after various time intervals, the cells were harvested and luciferase activity was determined as described above. As shown in Figure 3, a 100-fold induction of luciferase activity was observed after 5.5 hours incubation with Doxy. Fully induced levels were achieved in less than 24 hours of incubation with Doxy. Thus,

these results indicate that induction of gene expression occurs rapidly following exposure of the cells to the inducing agent.

EXAMPLE 3: Coordinate Regulation of the Expression of Two Nucleotide Sequences by a Tc-Controlled Transcriptional Activator

A recombinant expression vector for coordinate, bidirectional transcription of two nucleotide sequences was constructed comprising, in a 5' to 3' direction: a luciferase gene, a first minimal promoter, seven *tet* operator sequences, a second minimal promoter and a LacZ gene. The construct is illustrated in Figure 6. In this construct, the luciferase and LacZ genes are oriented such that they are transcribed in opposite orientations relative to the *tet* operator sequences, i.e., the luciferase gene is transcribed in a 5' to 3' direction from the bottom strand of DNA, whereas the LacZ gene is transcribed in a 5' to 3' direction from the top strand of DNA. The luciferase gene is followed by an SV40 polyadenylation signal, whereas the LacZ gene is followed by a β -globin polyadenylation signal.

The construct was transfected into the HeLa cell line HtTA-1 cells, which express a wild-type Tet repressor-VP16 fusion protein (referred to as tTA and described in Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551). The tTA fusion protein binds to *tet* operator sequences in the absence of Tc (or analogue) but not in the presence of Tc (or analogue). The construct was cotransfected into HtTA-1 cells with a plasmid which confers hygromycin resistance and stably transfected clones were selected based upon their hygromycin resistant phenotype. Selected hygromycin resistant (Hygr^r) clones were examined for luciferase and β -galactosidase activity. Clones positive for all three markers (Hygr^r, luc⁺, β -gal⁺) were then examined for tetracycline-dependent coregulation of expression of luciferase and β -galactosidase activity by culturing the clones in increasing amounts of tetracycline and measuring luciferase and β -galactosidase activity. The results of such an experiment using clone Ht1316-8/50 are shown in Figure 8. In the absence of tetracycline (in which case tTA can bind to tet operators and activate gene expression), both luciferase and β -galactosidase activity is detected. In the presence of increasing amounts of tetracycline, luciferase and β -galactosidase activity are coordinately and equivalently downregulated. This data demonstrates that expression of two genes can be coordinately regulated by a tetracycline-controlled transactivator by operatively linking the two genes to the same *tet* operator sequence(s).

EXAMPLE 4: Construction of a Tetracycline-Regulated Transcriptional Inhibitor Fusion Protein Comprising TetR and a Krueppel Silencer Domain

To construct an expression vector encoding a tetracycline-regulated transcriptional inhibitor of the invention (also referred to as a tetracycline controlled silencer domain, or

tSD), a nucleic acid fragment encoding a transcriptional silencer domain is ligated into an expression vector containing nucleotide sequences encoding a wild-type or modified (i.e., mutated) TetR such that the silencer domain coding sequences are ligated in-frame with the TetR coding sequences. The plasmid pUHD141sma-1 contains nucleotide sequences
5 encoding a wild-type Tn10-derived Tet repressor (the nucleotide and amino acid sequences of which are shown in SEQ ID NOs: 16 and 17, respectively). In pUHD141sma-1, the TetR coding sequence is linked at its 5' end to a CMV promoter and at its immediate 3' end to a nucleotide sequence that creates a polylinker into which additional nucleic acid fragments can be introduced. The nucleotide sequence across this polylinker region is: TCC CCG GGT
10 AAC TAA GTA AGG ATC C (SEQ ID NO: 24) (wherein TCC CCG GGT ACC encode amino acid residues 205-208 of TetR, namely Ser-Gly-Ser-Asn). This polylinker region includes restriction endonuclease sites for PspAI (CCC GGG) and BamHI (GGA TCC). Downstream of the polylinker region, the plasmid contains an SV40-derived polyadenylation signal. The pUHD141sma-1 vector is illustrated schematically in Figure 11.

15 To construct an expression vector encoding a fusion protein between TetR and a transcriptional silencer domain from the *Drosophila* Krueppel (Kr) protein, a nucleic acid fragment encoding a silencer domain from Kr is amplified by the polymerase chain reaction (PCR) using Kr cDNA as a template. Oligonucleotide primers are designed which amplify a nucleic acid fragment encoding the C-terminal 64 amino acids of Kr (referred to as C64KR).
20 This region corresponds to amino acid positions 403-466 of the native protein. The nucleotide and amino acid sequences of C64KR are shown in SEQ ID NO: 20 and SEQ ID NO: 21, respectively. PCR primers are designed to include restriction endonuclease sites such that the resultant amplified fragment contains restriction endonuclease sites at its 5' and 3' ends. Restriction endonuclease sites are chosen that are contained within the polylinker of
25 pUHD141sma-1 which allow in-frame, directional ligation of the amplified fragment into the polylinker site. For example, PCR primers are designed which incorporate a PspAI site (CCC GGG) at the 5' end of the fragment encoding C64KR and a BamHI site at the 3' end of the fragment. After a standard PCR reaction, the amplified fragment and pUHD141-sma1 are digested with PspAI and BamHI. The amplified fragment is then ligated directionally into
30 the polylinker site of pUHD141-sma1 using standard ligation conditions to create the expression vector pUHD141kr-1. Standard techniques are used to isolate the desired plasmid and confirm its construction. Construction of pUHD141kr-1 is illustrated schematically in Figure 11.

35 The resultant pUHD141kr-1 expression vector contains nucleotide sequences encoding a fusion protein comprising amino acids 1-207 of the wild type TetR linked in-frame to amino acids 403-466 of Kr (C64KR). The nucleotide and amino acid sequences across the junction of the fusion protein are as follows: AGT GGG TCC CCG GGT GAC ATG GAA (SEQ ID NO: 25) and Ser-Gly-Ser-Pro-Gly-Asp-Met-Glu (SEQ ID NO: 26). Ser-

Gly-Ser corresponds to amino acids 205-207 of TetR, Pro-Gly are encoded by the polylinker and Asp-Met-Glu correspond to the amino acids 403-405 of C64KR.

Similarly, an expression vector encoding a fusion protein of a mutated TetR (that binds to *tetO* only in the presence of Tc) and C64KR can be constructed as described above using nucleotide sequences encoding a mutant TetR (the nucleotide and amino acid sequences of which are shown in SEQ ID NOs: 18 and 19, respectively) in place of the wild type TetR sequences in pUHD141-sma1.

An expression vector encoding a TetR-Kr fusion protein (e.g., pUHD141kr-1) is transiently or stably transfected into host cells as described in Example 2 to express the TetR-Kr fusion protein in the host cell. A reporter gene construct containing one or more *tetO* sequences, a minimal promoter and a reporter gene, such as luciferase, is also transfected into the cells as described in Example 2. (Reporter gene constructs are described in further detail in USSN 08/076,726 and Gossen, M. and Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551). Luciferase activity in the presence and absence of increasing concentrations of Tc or an analogue, e.g., doxycycline, is measured as described in Example 2. For the wild type TetR-Kr fusion protein described above, the transcriptional inhibiting ability of the fusion protein is determined by comparing the amount of luciferase activity in the presence of doxycycline (no repression) to the amount of luciferase activity in the absence of doxycycline (repression). The transcriptional inhibiting activity of the fusion protein can also be tested using reporter gene constructs that exhibit higher basal levels of expression (i.e., higher levels of expression in the presence of doxycycline) by using a reporter gene construct that contains additional positive regulatory elements (e.g., enhancer sequences).

EXAMPLE 5: Construction of a Tetracycline-Regulated Transcriptional Inhibitor Fusion Protein Comprising TetR and a v-erbA Silencer Domain

To construct an expression vector encoding a fusion protein between TetR and a transcriptional silencer domain from the v-erbA oncogene product, a nucleic acid fragment encoding a silencer domain from v-erbA is ligated in-frame into pUHD141sma-1 as described in Example 4. A nucleic acid fragment encoding a v-erbA silencer domain suitable for ligation into pUHD141sma-1 is amplified by the polymerase chain reaction (PCR) using a v-erbA cDNA as a template. Oligonucleotide primers are designed which amplify a nucleic acid fragment encoding amino acids 364-635 of the native v-erbA protein. The nucleotide and amino acid sequences of this region of v-erbA are shown in SEQ ID NO: 22 and SEQ ID NO: 23, respectively. As described in Example 4, PCR primers are designed such that the amplified v-erbA fragment contains restriction endonuclease sites at its 5' and 3' ends, such as PspAI at the 5' end and BamHI at the 3' end. After a standard PCR reaction, the amplified fragment and pUHD141-sma1 are digested with PspAI and BamHI. The amplified fragment is then ligated directionally into the polylinker site of pUHD141-sma1 using standard ligation

conditions to create the expression vector pUHD141kr-1. Standard techniques can be used to isolate the desired plasmid and confirm its construction. Construction of pUHD141erb-1 is illustrated schematically in Figure 11.

5 The resultant pUHD141erb-1 expression vector contains nucleotide sequences encoding a fusion protein comprising a wild type TetR linked in-frame to amino acids 364-635 of v-erbA. The nucleotide and amino acid sequences across the junction of the fusion protein are as follows: AGT GGG TCC CCG GGT CTG GAC GAC (SEQ ID NO: 27) and Ser-Gly-Ser-Pro-Gly-Leu-Asp-Asp (SEQ ID NO: 28). Ser-Gly-Ser corresponds to amino acids 205-207 of TetR, Pro-Gly are encoded by the polylinker and Leu-Asp-Asp correspond
10 to amino acids 364-366 of the v-erbA silencer domain.

As described in Example 4, an expression vector encoding a fusion protein of a mutated TetR (that binds to *tetO* only in the presence of Tc) and a v-erbA silencer domain can be constructed as described above using nucleotide sequences encoding a mutant TetR (the nucleotide and amino acid sequences of which are shown in SEQ ID NOs: 18 and 19,
15 respectively) in place of the wild type TetR sequences in pUHD141-sma1.

Expression of the TetR-v-erbA fusion protein in host cells and assaying of the transcriptional inhibiting activity of the fusion protein is as described in Example 4 for the TetR-Kr fusion protein.

20 **EXAMPLE 6: Regulation of Gene Expression in Transgenic Animals by tTAR^R**

To examine the ability of tTAR^R to regulate gene expression *in vivo*, transgenic strains of mice were constructed which contained heterologous chromosomal insertions of either a tTAR^R expression construct or a reporter gene operably linked to *tet* operators. Single
25 transgenic strains containing either a tTAR^R expression construct or the *tetO*-linked reporter gene were then cross bred and double transgenic progeny were identified. The double transgenic animals were then characterized as to the ability of tTAR^R, in a tetracycline dependent manner, to regulate expression of the reporter gene. This example demonstrates that tTAR^R effectively stimulates the expression of a gene operably linked to *tet* operators in
30 tissues of the animals *in vivo* upon administration of tetracycline (or analogue) to the animals, whereas expression of the *tetO*-linked gene remains at background levels in the absence of tetracycline or an analogue. These results demonstrate that the tetracycline-controlled transcriptional regulatory system described herein functions effectively in animals, in addition to cell lines *in vitro*.

35

Generation of mice transgenic for a P_{hCMV}-tTAR^R expression unit

Mice expressing tTA protein were obtained by pronuclear injection into fertilized oocytes of a 2.7kb XhoI-Pfml fragment excised from plasmid pUHG17-1. This DNA fragment contained the tTAR^R gene (shown in SEQ ID NO: 1) under the transcriptional control

of the human CMV IE promoter (position +75 to -675) together with a rabbit β -globin polyadenylation site including an intron. The human CMV IE promoter is a constitutive promoter that allows expression of the mutated tetR-VP16 fusion protein in all cells where chromosomal integration of the DNA sequence encoding tTAR^R has occurred. DNA was
5 injected into fertilized oocytes at a concentration of approximately 5 ng per μ l by standard techniques. Transgenic mice were generated from the injected fertilized oocytes according to standard procedures. Transgenic founder mice were analyzed using polymerase chain reaction (PCR) and Southern hybridization to detect the presence of the tTAR^R transgene in
10 chromosomal DNA of the mice. Two transgenic mouse lines, CR3 and CR4 were identified and crossbred with another transgenic mouse line carrying a luciferase reporter gene under the control of *tetO* sequences. (described further below).

Generation of mice transgenic for the P_{hCMV}*-1 luciferase reporter unit

Mice carrying a P_{hCMV}*-1 luc reporter gene expression unit were generated by
15 pronuclear injection into fertilized oocytes of a 3.1 kb XhoI-EaeI fragment excised from plasmid pUHC13-3. This DNA-fragment contains the luciferase gene under transcriptional control of the tetracycline-responsive P_{hCMV}*-1 promoter (SEQ ID NO: 8), together with an SV40 t early polyadenylation site including an intron. DNA was injected into oocytes at a
20 concentration of approximately 5 ng per μ l and transgenic mice were generated according to standard procedures. Transgenic founder mice were analyzed using Southern hybridization to detect the presence of the P_{hCMV}*-1 luc transgene in chromosomal DNA of the mice. A mouse line transgenic for the *tetO*-linked luciferase reporter gene, L7, was crossbred with the
tTAR^R transgenic lines CR3 and CR4 (described further below).

25 Generation of mice transgenic for the P_{hCMV}*-1 luc and P_{hCMV} tTAR^R

Having constructed single transgenic mice expressing tTAR^R or carrying P_{hCMV}*-1 luc, double transgenic mice carrying both the tTA expression vector and the luciferase
reporter-units were obtained through cross breeding of heterozygous mice transgenic for one
of the two transgenes. Double transgenic animals were identified by standard screenings
30 (e.g., PCR and/or Southern hybridization) to detect the presence of both the tTAR^R transgene and the P_{hCMV}*-1 luc transgene in chromosomal DNA of the mice.

Induction and analysis of luciferase activity in tissue samples from mice

For oral administration, tetracycline or its derivative doxycycline were given in the
35 drinking water at a concentration of 200 μ g per ml with 5 % sucrose to hide the bitter taste of the antibiotics. For lactating mice, the concentration was 2 mg per ml with 10 % sucrose to ensure a sufficient uptake via the milk by the young.

To analyze luciferase activity, mice were killed by cervical dislocation and tissue samples were homogenized in 2 ml tubes containing 500 μ l lysis-buffer (25 mM Tris

phosphate, pH 7.8/2 mM DTT/ 2 mM EDTA/ 10 % glycerol/ 1 % Triton X100) using a Ultra-Turrax. The homogenate was frozen in liquid nitrogen and centrifuged after thawing for 5 min at 15,000g. 2-20 μ l of the supernatant were mixed with 250 μ l luciferase assay buffer (25 mM glycylglycine, pH 7.5/ 15 mM MgSO₄/ 5 mM ATP) and luciferase activity was measured for 10 sec after the injection of 100 μ l of a 125 μ M luciferin solution using Berthold Lumat LB 9501. The protein concentration of the homogenate was determined using Bradford assay and luciferase activity was calculated as relative light units (rlu) per μ g of total protein.

10 Results

Mice from 2 lines carrying the P_hCMV-tTAR^R transgene (CR3 and CR4) were mated with mice from line L7, transgenic for P_hCMV*-1 luc. The L7 line shows a very low but detectable background of luciferase activity in different organs that is probably due to position effects at the integration side. The background luciferase activity in different tissues of the L7 single transgenic mice is illustrated graphically in Figure 12, represented by the checked columns on the right-hand side for each tissue examined (each column represents the results from one animals). The luciferase activity in different tissues of the C3/L7 double transgenic mice in the absence of the tetracycline analogue doxycycline (i.e., uninduced conditions) is illustrated graphically in Figure 12, represented by the dark columns in the middle for each tissue examined. The luciferase activity in different tissues of the C3/L7 double transgenic mice in the presence of doxycycline (i.e., induced conditions) is illustrated graphically in Figure 12, represented by the light columns on the left-hand side for each tissue examined.

Luciferase activity was detectable in the seven tissues of the double transgenic mice examined: pancreas, kidney, stomach, muscle, thymus, heart and tongue. The tissue pattern of activated luciferase levels (i.e., in the presence of doxycycline) in the double transgenic mice was similar to expression patterns of the hCMV IE promoter reported in the literature. This is consistent with expression of the luciferase reporter gene being regulated by tTAR^R (which is expressed in the mice under the control of the hCMV IE promoter). The level of reporter gene induction varied among the different tissues examined. Regulation factors up to 100,000 fold (i.e., 5 orders of magnitude) were achieved, e.g. in the pancreas.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANTS:

10

- (A) NAME: Hermann Bujard
- (B) STREET: Remler Str. 9
- (C) CITY: D-69120 Heidelberg
- (E) COUNTRY: Germany
- (A) NAME: Manfred Gossen
- (B) STREET: 978 Arlington Boulevard #B
- (C) CITY: El Cerrito
- (D) STATE: California
- (E) COUNTRY: USA
- (F) POSTAL CODE (ZIP): 94530

15

- (ii) TITLE OF INVENTION: Tetracycline-Regulator Modulators

20

- (iii) NUMBER OF SEQUENCES: 28

(iv) CORRESPONDENCE ADDRESS:

25

- (A) ADDRESSEE: LAHIVE & COCKFIELD
- (B) STREET: 60 State Street, suite 510
- (C) CITY: Boston
- (D) STATE: Massachusetts
- (E) COUNTRY: USA
- (F) ZIP: 02109-1875

30

(v) COMPUTER READABLE FORM:

35

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: ASCII Text

(vii) PRIOR APPLICATION DATA:

40

- (A) APPLICATION NUMBER: To be assigned
- (B) FILING DATE: 07-JUN-1995
- (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA

45

- (A) APPLICATION NUMBER: US 08/383,754
- (B) FILING DATE: 03-FEB-1995
- (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA

50

- (A) APPLICATION NUMBER: US 08/275,876
- (B) FILING DATE: 15-JULY-94
- (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA

55

- (A) APPLICATION NUMBER: US 08/270,637
- (B) FILING DATE: 01-JULY-94
- (C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: DeConti, Giulio A. Jr.
- (B) REGISTRATION NUMBER: 31,503

- 68 -

(C) REFERENCE/DOCKET NUMBER: BBI-009C2PC

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617) 227-7400

(B) TELEFAX: (617) 227-5941

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1008 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: exon

(B) LOCATION: 1..1008

(ix) FEATURE:

(A) NAME/KEY: mRNA

(B) LOCATION: 1..1008

(ix) FEATURE:

(A) NAME/KEY: misc. binding

(B) LOCATION: 1..207

(ix) FEATURE:

(A) NAME/KEY: misc. binding

(B) LOCATION: 208..335

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1005

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

| | | |
|----|---|-----|
| 40 | ATG TCT AGA TTA GAT AAA AGT AAA GTG ATT AAC AGC GCA TTA GAG CTG | 48 |
| | Met Ser Arg Leu Asp Lys Ser Lys Val Ile Asn Ser Ala Leu Glu Leu | |
| | 1 5 10 15 | |
| 45 | CTT AAT GAG GTC GGA ATC GAA GGT TTA ACA ACC CGT AAA CTC GCC CAG | 96 |
| | Leu Asn Glu Val Gly Ile Glu Gly Leu Thr Thr Arg Lys Leu Ala Gln | |
| | 20 25 30 | |
| 50 | AAG CTA GGT GTA GAG CAG CCT ACA CTG TAT TGG CAT GTA AAA AAT AAG | 144 |
| | Lys Leu Gly Val Glu Gln Pro Thr Leu Tyr Trp His Val Lys Asn Lys | |
| | 35 40 45 | |
| | CGG GCT TTG CTC GAC GCC TTA GCC ATT GAG ATG TTA GAT AGG CAC CAT | 192 |
| | Arg Ala Leu Leu Asp Ala Leu Ala Ile Glu Met Leu Asp Arg His His | |
| | 50 55 60 | |

| | | |
|----|---|-----|
| | ACT CAC TTT TGC CCT TTA AAA GGG GAA AGC TGG CAA GAT TTT TTA CGC | 240 |
| | Thr His Phe Cys Pro Leu Lys Gly Glu Ser Trp Gln Asp Phe Leu Arg | |
| | 65 70 75 80 | |
| 5 | AAT AAG GCT AAA AGT TTT AGA TGT GCT TTA CTA AGT CAT CGC AAT GGA | 288 |
| | Asn Lys Ala Lys Ser Phe Arg Cys Ala Leu Leu Ser His Arg Asn Gly | |
| | 85 90 95 | |
| 10 | GCA AAA GTA CAT TCA GAT ACA CGG CCT ACA GAA AAA CAG TAT GAA ACT | 336 |
| | Ala Lys Val His Ser Asp Thr Arg Pro Thr Glu Lys Gln Tyr Glu Thr | |
| | 100 105 110 | |
| 15 | CTC GAA AAT CAA TTA GCC TTT TTA TGC CAA CAA GGT TTT TCA CTA GAG | 384 |
| | Leu Glu Asn Gln Leu Ala Phe Leu Cys Gln Gln Gly Phe Ser Leu Glu | |
| | 115 120 125 | |
| 20 | AAT GCA TTA TAT GCA CTC AGC GCT GTG GGG CAT TTT ACT TTA GGT TGC | 432 |
| | Asn Ala Leu Tyr Ala Leu Ser Ala Val Gly His Phe Thr Leu Gly Cys | |
| | 130 135 140 | |
| 25 | GTA TTG GAA GAT CAA GAG CAT CAA GTC GCT AAA GAA GAA AGG GAA ACA | 480 |
| | Val Leu Glu Asp Gln Glu His Gln Val Ala Lys Glu Glu Arg Glu Thr | |
| | 145 150 155 160 | |
| 30 | CCT ACT ACT GAT AGT ATG CCG CCA TTA TTA CGA CAA GCT ATC GAA TTA | 528 |
| | Pro Thr Thr Asp Ser Met Pro Pro Leu Leu Arg Gln Ala Ile Glu Leu | |
| | 165 170 175 | |
| 35 | TTT GAT CAC CAA GGT GCA GAG CCA GCC TTC TTA TTC GGC CTT GAA TTG | 576 |
| | Phe Asp His Gln Gly Ala Glu Pro Ala Phe Leu Phe Gly Leu Glu Leu | |
| | 180 185 190 | |
| 40 | ATC ATA TGC GGA TTA GAA AAA CAA CTT AAA TGT GAA AGT GGG TCC GCG | 624 |
| | Ile Ile Cys Gly Leu Glu Lys Gln Leu Lys Cys Glu Ser Gly Ser Ala | |
| | 195 200 205 | |
| 45 | TAC AGC CGC GCG CGT ACG AAA AAC AAT TAC GGG TCT ACC ATC GAG GGC | 672 |
| | Tyr Ser Arg Ala Arg Thr Lys Asn Asn Tyr Gly Ser Thr Ile Glu Gly | |
| | 210 215 220 | |
| 50 | CTG CTC GAT CTC CCG GAC GAC GAC GCC CCC GAA GAG GCG GGG CTG GCG | 720 |
| | Leu Leu Asp Leu Pro Asp Asp Asp Ala Pro Glu Glu Ala Gly Leu Ala | |
| | 225 230 235 240 | |
| 55 | GCT CCG CGC CTG TCC TTT CTC CCC GCG GGA CAC ACG CGC AGA CTG TCG | 768 |
| | Ala Pro Arg Leu Ser Phe Leu Pro Ala Gly His Thr Arg Arg Leu Ser | |
| | 245 250 255 | |
| 60 | ACG GCC CCC CCG ACC GAT GTC AGC CTG GGG GAC GAG CTC CAC TTA GAC | 816 |
| | Thr Ala Pro Pro Thr Asp Val Ser Leu Gly Asp Glu Leu His Leu Asp | |
| | 260 265 270 | |
| 65 | GGC GAG GAC GTG GCG ATG GCG CAT GCC GAC GCG CTA GAC GAT TTC GAT | 864 |
| | Gly Glu Asp Val Ala Met Ala His Ala Asp Ala Leu Asp Asp Phe Asp | |
| | 275 280 285 | |
| 70 | CTG GAC ATG TTG GGG GAC GGG GAT TCC CCG GGT CCG GGA TTT ACC CCC | 912 |
| | Leu Asp Met Leu Gly Asp Gly Asp Ser Pro Gly Pro Gly Phe Thr Pro | |
| | 290 295 300 | |

- 70 -

CAC GAC TCC GCC CCC TAC GGC GCT CTG GAT ATG GCC GAC TTC GAG TTT 960
 His Asp Ser Ala Pro Tyr Gly Ala Leu Asp Met Ala Asp Phe Glu Phe
 305 310 315 320

GAG CAG ATG TTT ACC GAT CCC CTT GGA ATT GAC GAG TAC GGT GGG TAG 1008
 Glu Gln Met Phe Thr Asp Pro Leu Gly Ile Asp Glu Tyr Gly Gly
 325 330 335

10 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 335 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ser Arg Leu Asp Lys Ser Lys Val Ile Asn Ser Ala Leu Glu Leu
 1 5 10 15

20 Leu Asn Glu Val Gly Ile Glu Gly Leu Thr Thr Arg Lys Leu Ala Gln
 20 25 30

Lys Leu Gly Val Glu Gln Pro Thr Leu Tyr Trp His Val Lys Asn Lys
 35 40 45

25 Arg Ala Leu Leu Asp Ala Leu Ala Ile Glu Met Leu Asp Arg His His
 50 55 60

30 Thr His Phe Cys Pro Leu Lys Gly Glu Ser Trp Gln Asp Phe Leu Arg
 65 70 75 80

Asn Lys Ala Lys Ser Phe Arg Cys Ala Leu Leu Ser His Arg Asn Gly
 85 90 95

35 Ala Lys Val His Ser Asp Thr Arg Pro Thr Glu Lys Gln Tyr Glu Thr
 100 105 110

Leu Glu Asn Gln Leu Ala Phe Leu Cys Gln Gln Gly Phe Ser Leu Glu
 115 120 125

40 Asn Ala Leu Tyr Ala Leu Ser Ala Val Gly His Phe Thr Leu Gly Cys
 130 135 140

45 Val Leu Glu Asp Gln Glu His Gln Val Ala Lys Glu Glu Arg Glu Thr
 145 150 155 160

Pro Thr Thr Asp Ser Met Pro Pro Leu Leu Arg Gln Ala Ile Glu Leu
 165 170 175

50 Phe Asp His Gln Gly Ala Glu Pro Ala Phe Leu Phe Gly Leu Glu Leu
 180 185 190

Ile Ile Cys Gly Leu Glu Lys Gln Leu Lys Cys Glu Ser Gly Ser Ala
 195 200 205

55

- 71 -

Tyr Ser Arg Ala Arg Thr Lys Asn Asn Tyr Gly Ser Thr Ile Glu Gly
 210 215 220
 5 Leu Leu Asp Leu Pro Asp Asp Ala Pro Glu Glu Ala Gly Leu Ala
 225 230 235 240
 Ala Pro Arg Leu Ser Phe Leu Pro Ala Gly His Thr Arg Arg Leu Ser
 245 250 255
 10 Thr Ala Pro Pro Thr Asp Val Ser Leu Gly Asp Glu Leu His Leu Asp
 260 265 270
 Gly Glu Asp Val Ala Met Ala His Ala Asp Ala Leu Asp Asp Phe Asp
 275 280 285
 15 Leu Asp Met Leu Gly Asp Gly Asp Ser Pro Gly Pro Gly Phe Thr Pro
 290 295 300
 20 His Asp Ser Ala Pro Tyr Gly Ala Leu Asp Met Ala Asp Phe Glu Phe
 305 310 315 320
 Glu Gln Met Phe Thr Asp Pro Leu Gly Ile Asp Glu Tyr Gly Gly
 325 330 335
 25

INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
 30
 (ii) MOLECULE TYPE: DNA
 35
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAC GCG CTA GAC GAT TTC GAT CTG GAC ATG TTG 33
 Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu
 40 1 5 10

(2) INFORMATION FOR SEQ ID NO:4:
 45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 50 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Pro Lys Arg Pro Arg Pro
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 569 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GAATTCGGGG CCGCGGAGGC TGGATCGGTC CCGGTGTCTT CTATGGAGGT CAAAACAGCG 60
TGGATGGCGT CTCCAGGCGA TCTGACGGTT CACTAAACGA GCTCTGCTTA TATAGGTCGA 120
GTTTACCACT CCCTATCAGT GATAGAGAAA AGTGAAAGTC GAGTTTACCA CTCCCTATCA 180
GTGATAGAGA AAAGTGAAAG TCGAGTTTAC CACTCCCTAT CAGTGATAGA GAAAAGTGAA 240
AGTCGAGTTT ACCACTCCCT ACCAGTGATA GAGAAAAGTG AAAGTCGAGT TTACCACTCC 300
CTATCAGTGA TAGAGAAAAG TGAAAGTCGA GTTTACCACT CCCTATCAGT GATAGAGAAA 360
AGTGAAAGTC GAGTTTACCA CTCCCTATCA GTGATAGAGA AAAGTGAAAG TCGAGCTCGG 420
TACCCGGGTC GAGTAGGCGT GTACGGTGGG AGGCCTATAT AAGCAGAGCT CGTTTAGTGA 480
ACCGTCAGAT CGCCTGGAGA CGCCATCCAC GCTGTTTTGA CCTCCATAGA AGACACCGGG 540
ACCGATCCAG CCTCCGCGGC CCCGAATTC 569

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 520 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AGATCTGCAG GGTCTGCTCGG TGTTCGAGGC CACACGCGTC ACCTTAATAT GCGAAGTGGA 60
CCGGATCTCG AGTTTACCAC TCCCTATCAG TGATAGAGAA AAGTGAAAGT CGAGTTTACC 120
ACTCCCTATC AGTGATAGAG AAAAGTGAAA GTCGAGTTTA CCACTCCCTA TCAGTGATAG 180
AGAAAAGTGA AAGTCGAGTT TACCACTCCC TATCAGTGAT AGAGAAAAGT GAAAGTCGAG 240
TTTACCACTC CCTATCAGTG ATAGAGAAAA GTGAAAGTCG AGTTTACCAC TCCCTATCAG 300
TGATAGAGAA AAGTGAAAGT CGAGTTTACC ACTCCCTATC AGTGATAGAG AAAAGTGAAA 360
GTCGAGCTCG GTACCCGGGT CGAGTAGGCG TGTACGGTGG GAGGCCTATA TAAGCAGAGC 420
TCGTTTAGTG AACCGTCAGA TCGCCTGGAG ACGCCATCCA CGCTGTTTTG ACCTCCATAG 480
AAGACACCGG GACCGATCCA GCCTCCGCGG CCCC GAATTC 520

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 450 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Human cytomegalovirus
(B) STRAIN: K12, Towne

(ix) FEATURE:

- (A) NAME/KEY: mRNA
(B) LOCATION: 382..450

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GAATTCCTCG AGTTTACCAC TCCCTATCAG TGATAGAGAA AAGTGAAAGT CGAGTTTACC 60
ACTCCCTATC AGTGATAGAG AAAAGTGAAA GTCGAGTTTA CCACTCCCTA TCAGTGATAG 120
AGAAAAGTGA AAGTCGAGTT TACCACTCCC TATCAGTGAT AGAGAAAAGT GAAAGTCGAG 180
TTTACCACTC CCTATCAGTG ATAGAGAAAA GTGAAAGTCG AGTTTACCAC TCCCTATCAG 240

- 74 -

TGATAGAGAA AAGTGAAAGT CGAGTTTACC ACTCCCTATC AGTGATAGAG AAAAGTGAAA 300
GTCGAGCTCG GTACCCGGGT CGAGTAGGCG TGTACGGTGG GAGGCCTATA TAAGCAGAGC 360
TCGTTTAGTG AACCGTCAGA TCGCCTGGAG ACGCCATCCA CGCTGTTTTG ACCTCCATAG 420
AAGACACCGG GACCGATCCA GCCTCCGCGG 450

5

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 450 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Human cytomegalovirus
(B) STRAIN: Towne

20

(ix) FEATURE:

- (A) NAME/KEY: mRNA
(B) LOCATION: 382..450

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GAATTCCTCG ACCCGGGTAC CGAGCTCGAC TTTCACTTTT CTCTATCACT GATAGGGAGT 60
25 GGTAAGTCTCG ACTTTCACCT TTCTCTATCA CTGATAGGGA GTGGTAAACT CGACTTTCAC 120
TTTTCTCTAT CACTGATAGG GAGTGGTAAA CTCGACTTTC ACTTTTCTCT ATCACTGATA 180
GGGAGTGGTA AACTCGACTT TCACTTTTCT CTATCACTGA TAGGGAGTGG TAAACTCGAC 240
TTTCACTTTT CTCTATCACT GATAGGGAGT GGTAAGTCTCG ACTTTCACCT TTCTCTATCA 300
CTGATAGGGA GTGGTAAACT CGAGTAGGCG TGTACGGTGG GAGGCCTATA TAAGCAGAGC 360
30 TCGTTTAGTG AACCGTCAGA TCGCCTGGAG ACGCCATCCA CGCTGTTTTG ACCTCCATAG 420
AAGACACCGG GACCGATCCA GCCTCCGCGG 450

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 398 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

40

- 75 -

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Herpes Simplex Virus

(B) STRAIN: KOS

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GAGCTCGACT TTCAC TTTTC TCTATCACTG ATAGGGAGTG GTAAACTCGA CTTTCACTTT 60
TCTCTATCAC TGATAGGGAG TGGTAAACTC GACTTTCACT TTTCTCTATC ACTGATAGGG 120
AGTGGTAAAC TCGACTTTCA CTTTTCTCTA TCACTGATAG GGAGTGGTAA ACTCGACTTT 180
CACTTTTCTC TATCACTGAT AGGGAGTGGT AACTCGACT TTCAC TTTTC TCTATCACTG 240
10 ATAGGGAGTG GTAAACTCGA CTTTCACTTT TCTCTATCAC TGATAGGGAG TGGTAAACTC 300
GAGATCCGGC GAATTCGAAC ACGCAGATGC AGTCGGGGCG GCGCGGTCCG AGGTCCACTT 360
CGCATATTAA GGTGACGCGT GTGGCCTCGA ACACCGAG 398

(2) INFORMATION FOR SEQ ID NO:11:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ACTTTATCAC TGATAAACAA ACTTATCAGT GATAAAGA 38

25 (2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ACTCTATCAT TGATAGAGTT CCCTATCAGT GATAGAGA 38

35

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AGCTTATCAT CGATAAGCTA GTTTATCACA GTTAAATT

38

10

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

20 ACTCTATCAT TGATAGGGAA CTCTATCAAT GATAGGGA

38

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AATCTATCAC TGATAGAGTA CCCTATCATC GATAGAGA

38

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 621 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

35

40 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

- 77 -

| | | |
|----|---|-----|
| | ATG TCT AGA TTA GAT AAA AGT AAA GTG ATT AAC AGC GCA TTA GAG CTG | 48 |
| | Met Ser Arg Leu Asp Lys Ser Lys Val Ile Asn Ser Ala Leu Glu Leu | |
| | 1 5 10 15 | |
| 5 | CTT AAT GAG GTC GGA ATC GAA GGT TTA ACA ACC CGT AAA CTC GCC CAG | 96 |
| | Leu Asn Glu Val Gly Ile Glu Gly Leu Thr Thr Arg Lys Leu Ala Gln | |
| | 20 25 30 | |
| 10 | AAG CTA GGT GTA GAG CAG CCT ACA TTG TAT TGG CAT GTA AAA AAT AAG | 144 |
| | Lys Leu Gly Val Glu Gln Pro Thr Leu Tyr Trp His Val Lys Asn Lys | |
| | 35 40 45 | |
| 15 | CGG GCT TTG CTC GAC GCC TTA GCC ATT GAG ATG TTA GAT AGG CAC CAT | 192 |
| | Arg Ala Leu Leu Asp Ala Leu Ala Ile Glu Met Leu Asp Arg His His | |
| | 50 55 60 | |
| 20 | ACT CAC TTT TGC CCT TTA GAA GGG GAA AGC TGG CAA GAT TTT TTA CGT | 240 |
| | Thr His Phe Cys Pro Leu Glu Gly Glu Ser Trp Gln Asp Phe Leu Arg | |
| | 65 70 75 80 | |
| 25 | AAT AAG GCT AAA AGT TTT AGA TGT GCT TTA CTA AGT CAT CGC GAT GGA | 288 |
| | Asn Lys Ala Lys Ser Phe Arg Cys Ala Leu Leu Ser His Arg Asp Gly | |
| | 85 90 95 | |
| 30 | GCA AAA GTA CAT TTA GGT ACA CGG CCT ACA GAA AAA CAG TAT GAA ACT | 336 |
| | Ala Lys Val His Leu Gly Thr Arg Pro Thr Glu Lys Gln Tyr Glu Thr | |
| | 100 105 110 | |
| 35 | CTC GAA AAT CAA TTA GCC TTT TTA TGC CAA CAA GGT TTT TCA CTA GAG | 384 |
| | Leu Glu Asn Gln Leu Ala Phe Leu Cys Gln Gln Gly Phe Ser Leu Glu | |
| | 115 120 125 | |
| 40 | AAT GCA TTA TAT GCA CTC AGC GCT GTG GGG CAT TTT ACT TTA GGT TGC | 432 |
| | Asn Ala Leu Tyr Ala Leu Ser Ala Val Gly His Phe Thr Leu Gly Cys | |
| | 130 135 140 | |
| 45 | GTA TTG GAA GAT CAA GAG CAT CAA GTC GCT AAA GAA GAA AGG GAA ACA | 480 |
| | Val Leu Glu Asp Gln Glu His Gln Val Ala Lys Glu Glu Arg Glu Thr | |
| | 145 150 155 160 | |
| 50 | CCT ACT ACT GAT AGT ATG CCG CCA TTA TTA CGA CAA GCT ATC GAA TTA | 528 |
| | Pro Thr Thr Asp Ser Met Pro Pro Leu Leu Arg Gln Ala Ile Glu Leu | |
| | 165 170 175 | |
| 55 | TTT GAT CAC CAA GGT GCA GAG CCA GCC TTC TTA TTC GGC CTT GAA TTG | 576 |
| | Phe Asp His Gln Gly Ala Glu Pro Ala Phe Leu Phe Gly Leu Glu Leu | |
| | 180 185 190 | |
| 60 | ATC ATA TGC GGA TTA GAA AAA CAA CTT AAA TGT GAA AGT GGG TCC | 621 |
| | Ile Ile Cys Gly Leu Glu Lys Gln Leu Lys Cys Glu Ser Gly Ser | |
| | 195 200 205 | |

(2) INFORMATION FOR SEQ ID NO:17:

- 55 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 207 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

5 Met Ser Arg Leu Asp Lys Ser Lys Val Ile Asn Ser Ala Leu Glu Leu
 1 5 10 15
 Leu Asn Glu Val Gly Ile Glu Gly Leu Thr Thr Arg Lys Leu Ala Gln
 20 25 30
 10 Lys Leu Gly Val Glu Gln Pro Thr Leu Tyr Trp His Val Lys Asn Lys
 35 40 45
 Arg Ala Leu Leu Asp Ala Leu Ala Ile Glu Met Leu Asp Arg His His
 50 55 60
 15 Thr His Phe Cys Pro Leu Glu Gly Glu Ser Trp Gln Asp Phe Leu Arg
 65 70 75 80
 Asn Lys Ala Lys Ser Phe Arg Cys Ala Leu Leu Ser His Arg Asp Gly
 85 90 95
 20 Ala Lys Val His Leu Gly Thr Arg Pro Thr Glu Lys Gln Tyr Glu Thr
 100 105 110
 Leu Glu Asn Gln Leu Ala Phe Leu Cys Gln Gln Gly Phe Ser Leu Glu
 115 120 125
 Asn Ala Leu Tyr Ala Leu Ser Ala Val Gly His Phe Thr Leu Gly Cys
 130 135 140
 30 Val Leu Glu Asp Gln Glu His Gln Val Ala Lys Glu Glu Arg Glu Thr
 145 150 155 160
 Pro Thr Thr Asp Ser Met Pro Pro Leu Leu Arg Gln Ala Ile Glu Leu
 165 170 175
 35 Phe Asp His Gln Gly Ala Glu Pro Ala Phe Leu Phe Gly Leu Glu Leu
 180 185 190
 Ile Ile Cys Gly Leu Glu Lys Gln Leu Lys Cys Glu Ser Gly Ser
 195 200 205
 40

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 621 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

- 79 -

| | | |
|----|---|-----|
| | ATG TCT AGA TTA GAT AAA AGT AAA GTG ATT AAC AGC GCA TTA GAG CTG | 48 |
| | Met Ser Arg Leu Asp Lys Ser Lys Val Ile Asn Ser Ala Leu Glu Leu | |
| | 1 5 10 15 | |
| 5 | CTT AAT GAG GTC GGA ATC GAA GGT TTA ACA ACC CGT AAA CTC GCC CAG | 96 |
| | Leu Asn Glu Val Gly Ile Glu Gly Leu Thr Thr Arg Lys Leu Ala Gln | |
| | 20 25 30 | |
| 10 | AAG CTA GGT GTA GAG CAG CCT ACA CTG TAT TGG CAT GTA AAA AAT AAG | 144 |
| | Lys Leu Gly Val Glu Gln Pro Thr Leu Tyr Trp His Val Lys Asn Lys | |
| | 35 40 45 | |
| 15 | CGG GCT TTG CTC GAC GCC TTA GCC ATT GAG ATG TTA GAT AGG CAC CAT | 192 |
| | Arg Ala Leu Leu Asp Ala Leu Ala Ile Glu Met Leu Asp Arg His His | |
| | 50 55 60 | |
| 20 | ACT CAC TTT TGC CCT TTA AAA GGG GAA AGC TGG CAA GAT TTT TTA CGC | 240 |
| | Thr His Phe Cys Pro Leu Lys Gly Glu Ser Trp Gln Asp Phe Leu Arg | |
| | 65 70 75 80 | |
| 25 | AAT AAG GCT AAA AGT TTT AGA TGT GCT TTA CTA AGT CAT CGC AAT GGA | 288 |
| | Asn Lys Ala Lys Ser Phe Arg Cys Ala Leu Leu Ser His Arg Asn Gly | |
| | 85 90 95 | |
| 30 | GCA AAA GTA CAT TCA GAT ACA CGG CCT ACA GAA AAA CAG TAT GAA ACT | 336 |
| | Ala Lys Val His Ser Asp Thr Arg Pro Thr Glu Lys Gln Tyr Glu Thr | |
| | 100 105 110 | |
| 35 | CTC GAA AAT CAA TTA GCC TTT TTA TGC CAA CAA GGT TTT TCA CTA GAG | 384 |
| | Leu Glu Asn Gln Leu Ala Phe Leu Cys Gln Gln Gly Phe Ser Leu Glu | |
| | 115 120 125 | |
| 40 | AAT GCA TTA TAT GCA CTC AGC GCT GTG GGG CAT TTT ACT TTA GGT TGC | 432 |
| | Asn Ala Leu Tyr Ala Leu Ser Ala Val Gly His Phe Thr Leu Gly Cys | |
| | 130 135 140 | |
| 45 | GTA TTG GAA GAT CAA GAG CAT CAA GTC GCT AAA GAA GAA AGG GAA ACA | 480 |
| | Val Leu Glu Asp Gln Glu His Gln Val Ala Lys Glu Glu Arg Glu Thr | |
| | 145 150 155 160 | |
| 50 | CCT ACT ACT GAT AGT ATG CCG CCA TTA TTA CGA CAA GCT ATC GAA TTA | 528 |
| | Pro Thr Thr Asp Ser Met Pro Pro Leu Leu Arg Gln Ala Ile Glu Leu | |
| | 165 170 175 | |
| 55 | TTT GAT CAC CAA GGT GCA GAG CCA GCC TTC TTA TTC GGC CTT GAA TTG | 576 |
| | Phe Asp His Gln Gly Ala Glu Pro Ala Phe Leu Phe Gly Leu Glu Leu | |
| | 180 185 190 | |
| 60 | ATC ATA TGC GGA TTA GAA AAA CAA CTT AAA TGT GAA AGT GGG TCC | 621 |
| | Ile Ile Cys Gly Leu Glu Lys Gln Leu Lys Cys Glu Ser Gly Ser | |
| | 195 200 205 | |

(2) INFORMATION FOR SEQ ID NO:19:

- 55 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 207 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

- 80 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

5 Met Ser Arg Leu Asp Lys Ser Lys Val Ile Asn Ser Ala Leu Glu Leu
 1 5 10 15
 Leu Asn Glu Val Gly Ile Glu Gly Leu Thr Thr Arg Lys Leu Ala Gln
 20 25 30
 10 Lys Leu Gly Val Glu Gln Pro Thr Leu Tyr Trp His Val Lys Asn Lys
 35 40 45
 Arg Ala Leu Leu Asp Ala Leu Ala Ile Glu Met Leu Asp Arg His His
 50 55 60
 15 Thr His Phe Cys Pro Leu Lys Gly Glu Ser Trp Gln Asp Phe Leu Arg
 65 70 75 80
 Asn Lys Ala Lys Ser Phe Arg Cys Ala Leu Leu Ser His Arg Asn Gly
 85 90 95
 20 Ala Lys Val His Ser Asp Thr Arg Pro Thr Glu Lys Gln Tyr Glu Thr
 100 105 110
 Leu Glu Asn Gln Leu Ala Phe Leu Cys Gln Gln Gly Phe Ser Leu Glu
 115 120 125
 Asn Ala Leu Tyr Ala Leu Ser Ala Val Gly His Phe Thr Leu Gly Cys
 130 135 140
 30 Val Leu Glu Asp Gln Glu His Gln Val Ala Lys Glu Glu Arg Glu Thr
 145 150 155 160
 Pro Thr Thr Asp Ser Met Pro Pro Leu Leu Arg Gln Ala Ile Glu Leu
 165 170 175
 35 Phe Asp His Gln Gly Ala Glu Pro Ala Phe Leu Phe Gly Leu Glu Leu
 180 185 190
 Ile Ile Cys Gly Leu Glu Lys Gln Leu Lys Cys Glu Ser Gly Ser
 195 200 205
 40

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 192 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

- 81 -

```

      GAC ATG GAA AAA GCG ACA CCG GAG ACG ATG GTC CAT TGG ATT TGT CTG      48
      Asp Met Glu Lys Ala Thr Pro Glu Thr Met Val His Trp Ile Cys Leu
      1              5              10              15

5     AAG ATG GAG CCA GCT CTG TGG ATG GCC ATT ACA GCA ACA TCG CAC GGC      96
      Lys Met Glu Pro Ala Leu Trp Met Ala Ile Thr Ala Thr Ser His Gly
              20              25              30

10    GCA AGG CAC AGG ACA TTC GTC GGG TTT TCC GGC TGC CTC CAC CGC AAA      144
      Ala Arg His Arg Thr Phe Val Gly Phe Ser Gly Cys Leu His Arg Lys
              35              40              45

      TCC CTC ACG TAC CCA GTG ATA TGC CTG AGC AAA CCG AGC CAG AGG ATT      192
      Ser Leu Thr Tyr Pro Val Ile Cys Leu Ser Lys Pro Ser Gln Arg Ile
15    50              55              60

```

(2) INFORMATION FOR SEQ ID NO:21:

- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 64 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```

      Asp Met Glu Lys Ala Thr Pro Glu Thr Met Val His Trp Ile Cys Leu
      1              5              10              15

30    Lys Met Glu Pro Ala Leu Trp Met Ala Ile Thr Ala Thr Ser His Gly
              20              25              30

      Ala Arg His Arg Thr Phe Val Gly Phe Ser Gly Cys Leu His Arg Lys
              35              40              45

35    Ser Leu Thr Tyr Pro Val Ile Cys Leu Ser Lys Pro Ser Gln Arg Ile
      50              55              60

```

(2) INFORMATION FOR SEQ ID NO:22:

- 40 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 816 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- 45 (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

50    CTG GAC GAC TCG AAG CGC GTA GCC AAG CGG AAG CTG ATC GAG GAG AAC      48
      Leu Asp Asp Ser Lys Arg Val Ala Lys Arg Lys Leu Ile Glu Glu Asn
      1              5              10              15

```

| | | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | CGG | GAG | CGG | CGA | CGC | AAG | GAG | GAG | ATG | ATC | AAA | TCC | CTG | CAG | CAC | CGG | 96 |
| | Arg | Glu | Arg | Arg | Arg | Lys | Glu | Glu | Met | Ile | Lys | Ser | Leu | Gln | His | Arg | |
| | | | | 20 | | | | | 25 | | | | | 30 | | | |
| 5 | CCC | AGC | CCC | AGC | GCA | GAG | GAG | TGG | GAG | CTG | ATC | CAC | GTG | GTG | ACC | GAG | 144 |
| | Pro | Ser | Pro | Ser | Ala | Glu | Glu | Trp | Glu | Leu | Ile | His | Val | Val | Thr | Glu | |
| | | | 35 | | | | | 40 | | | | | 45 | | | | |
| 10 | GCG | CAC | CGC | AGC | ACC | AAC | GCG | CAG | GGC | AGC | CAC | TGG | AAG | CAG | AGG | AGG | 192 |
| | Ala | His | Arg | Ser | Thr | Asn | Ala | Gln | Gly | Ser | His | Trp | Lys | Gln | Arg | Arg | |
| | | 50 | | | | | 55 | | | | | 60 | | | | | |
| 15 | AAA | TTC | CTG | CTC | GAA | GAT | ATC | GGT | CAG | TCG | CCC | ATG | GCC | TCC | ATG | CTT | 240 |
| | Lys | Phe | Leu | Leu | Glu | Asp | Ile | Gly | Gln | Ser | Pro | Met | Ala | Ser | Met | Leu | |
| | 65 | | | | | 70 | | | | | 75 | | | | | 80 | |
| 20 | GAC | GGG | GAC | AAA | GTG | GAC | CTG | GAG | GCG | TTC | AGC | GAG | TTT | ACA | AAA | ATC | 288 |
| | Asp | Gly | Asp | Lys | Val | Asp | Leu | Glu | Ala | Phe | Ser | Glu | Phe | Thr | Lys | Ile | |
| | | | | 85 | | | | | | 90 | | | | | 95 | | |
| 25 | ATC | ACG | CCG | GCC | ATC | ACC | CGC | GTG | GTC | GAC | TTT | GCC | AAA | AAC | CTG | CCC | 336 |
| | Ile | Thr | Pro | Ala | Ile | Thr | Arg | Val | Val | Asp | Phe | Ala | Lys | Asn | Leu | Pro | |
| | | | | 100 | | | | | 105 | | | | | 110 | | | |
| 30 | ATG | TTC | TCG | GAG | CTG | CCG | TGC | GAG | GAT | CAG | ATC | ATC | CTG | CTG | AAG | GGC | 384 |
| | Met | Phe | Ser | Glu | Leu | Pro | Cys | Glu | Asp | Gln | Ile | Ile | Leu | Leu | Lys | Gly | |
| | | | 115 | | | | | 120 | | | | | 125 | | | | |
| 35 | TGC | TGC | ATG | GAG | ATC | ATG | TCG | CTG | CGC | GCC | GCC | GTG | CGC | TAC | GAC | CCC | 432 |
| | Cys | Cys | Met | Glu | Ile | Met | Ser | Leu | Arg | Ala | Ala | Val | Arg | Tyr | Asp | Pro | |
| | | | 130 | | | | | 135 | | | | | 140 | | | | |
| 40 | GAG | AGC | GAA | ACG | CTG | ACG | CTG | AGC | GGG | GAA | ATG | GCC | GTC | AAA | CGC | GAG | 480 |
| | Glu | Ser | Glu | Thr | Leu | Thr | Leu | Ser | Gly | Glu | Met | Ala | Val | Lys | Arg | Glu | |
| | 145 | | | | | 150 | | | | | 155 | | | | | 160 | |
| 45 | CAG | TTG | AAG | AAC | GGA | GGG | CTG | GGG | GTC | GTG | TCT | GAT | GCC | ATC | TTC | GAC | 528 |
| | Gln | Leu | Lys | Asn | Gly | Gly | Leu | Gly | Val | Val | Ser | Asp | Ala | Ile | Phe | Asp | |
| | | | | 165 | | | | | 170 | | | | | | 175 | | |
| 50 | CTC | GGC | AAG | TCG | CTG | TCT | GCC | TTC | AAC | CTG | GAC | GAC | ACC | GAG | GTG | GCC | 576 |
| | Leu | Gly | Lys | Ser | Leu | Ser | Ala | Phe | Asn | Leu | Asp | Asp | Thr | Glu | Val | Ala | |
| | | | 180 | | | | | | 185 | | | | | 190 | | | |
| 55 | CTG | CTG | CAG | GCC | GTG | CTG | CTC | ATG | TCC | TCA | GAC | CGG | ACG | GGG | CTG | ATC | 624 |
| | Leu | Leu | Gln | Ala | Val | Leu | Leu | Met | Ser | Ser | Asp | Arg | Thr | Gly | Leu | Ile | |
| | | | 195 | | | | | 200 | | | | | 205 | | | | |
| 60 | TGC | GTG | GAT | AAG | ATA | GAG | AAG | TGC | CAG | GAG | TCG | TAC | CTG | CTG | GCG | TTC | 672 |
| | Cys | Val | Asp | Lys | Ile | Glu | Lys | Cys | Gln | Glu | Ser | Tyr | Leu | Leu | Ala | Phe | |
| | | 210 | | | | | | 215 | | | | | 220 | | | | |
| 65 | GAG | CAC | TAC | ATC | AAC | TAC | CGC | AAA | CAC | AAC | ATT | CCC | CAC | TTC | TGG | TCC | 720 |
| | Glu | His | Tyr | Ile | Asn | Tyr | Arg | Lys | His | Asn | Ile | Pro | His | Phe | Trp | Ser | |
| | 225 | | | | | 230 | | | | | 235 | | | | | 240 | |

- 83 -

AAG CTG CTG ATG AAG GTG GCG GAC CTG CGC ATG ATC GGC GCC TAC CAC 768
 Lys Leu Leu Met Lys Val Ala Asp Leu Arg Met Ile Gly Ala Tyr His
 245 250 255

5 GCC AGC CGC TTC CTG CAC ATG AAG GTG GAG TGC CCC ACC GAG CTC TCC 816
 Ala Ser Arg Phe Leu His Met Lys Val Glu Cys Pro Thr Glu Leu Ser
 260 265 270

10 (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 272 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Leu Asp Asp Ser Lys Arg Val Ala Lys Arg Lys Leu Ile Glu Glu Asn
 1 5 10 15
 20 Arg Glu Arg Arg Arg Lys Glu Glu Met Ile Lys Ser Leu Gln His Arg
 20 25 30
 25 Pro Ser Pro Ser Ala Glu Glu Trp Glu Leu Ile His Val Val Thr Glu
 35 40 45
 Ala His Arg Ser Thr Asn Ala Gln Gly Ser His Trp Lys Gln Arg Arg
 50 55 60
 30 Lys Phe Leu Leu Glu Asp Ile Gly Gln Ser Pro Met Ala Ser Met Leu
 65 70 75 80
 Asp Gly Asp Lys Val Asp Leu Glu Ala Phe Ser Glu Phe Thr Lys Ile
 85 90 95
 35 Ile Thr Pro Ala Ile Thr Arg Val Val Asp Phe Ala Lys Asn Leu Pro
 100 105 110
 40 Met Phe Ser Glu Leu Pro Cys Glu Asp Gln Ile Ile Leu Leu Lys Gly
 115 120 125
 Cys Cys Met Glu Ile Met Ser Leu Arg Ala Ala Val Arg Tyr Asp Pro
 130 135 140
 45 Glu Ser Glu Thr Leu Thr Leu Ser Gly Glu Met Ala Val Lys Arg Glu
 145 150 155 160
 Gln Leu Lys Asn Gly Gly Leu Gly Val Val Ser Asp Ala Ile Phe Asp
 165 170 175
 50 Leu Gly Lys Ser Leu Ser Ala Phe Asn Leu Asp Asp Thr Glu Val Ala
 180 185 190
 55 Leu Leu Gln Ala Val Leu Leu Met Ser Ser Asp Arg Thr Gly Leu Ile
 195 200 205

Cys Val Asp Lys Ile Glu Lys Cys Gln Glu Ser Tyr Leu Leu Ala Phe
210 215 220

5 Glu His Tyr Ile Asn Tyr Arg Lys His Asn Ile Pro His Phe Trp Ser
225 230 235 240

Lys Leu Leu Met Lys Val Ala Asp Leu Arg Met Ile Gly Ala Tyr His
245 250 255

10 Ala Ser Arg Phe Leu His Met Lys Val Glu Cys Pro Thr Glu Leu Ser
260 265 270

(2) INFORMATION FOR SEQ ID NO:24:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

- 20 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

TCCCCGGGTA ACTAAGTAAG GATCC

25

25 (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
30 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

35 AGTGGGTCCC CGGGTGACAT GGAA

24

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
40 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: polypeptide

- (v) FRAGMENT TYPE: internal

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

- 85 -

Ser Gly Ser Pro Gly Asp Met Glu
1 5

(2) INFORMATION FOR SEQ ID NO:27:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

- 10 (ii) MOLECULE TYPE: DNA

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

AGTGGGTCCC CGGGTCTGGA CGAC

24

15

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
20 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: polypeptide

- (v) FRAGMENT TYPE: internal

- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Ser Gly Ser Pro Gly Leu Asp Asp

1

5

CLAIMS

1. An isolated nucleic acid encoding a fusion protein which activates transcription, the fusion protein comprising a first polypeptide which binds to a *tet* operator sequence in the presence of tetracycline or a tetracycline analogue operatively linked to a second polypeptide which activates transcription in eukaryotic cells.
2. The nucleic acid of claim 1, wherein the first polypeptide is a mutated Tet repressor.
3. The nucleic acid of claim 2, wherein the mutated Tet repressor has at least one amino acid substitution compared to a wild-type Tet repressor.
4. The nucleic acid of claim 2, wherein the mutated Tet repressor has at least one amino acid addition or deletion compared to a wild-type Tet repressor.
5. The nucleic acid of claim 3, wherein the mutated Tet repressor is a mutated Tn10-derived Tet repressor having an amino acid substitution at at least one amino acid position selected from the group consisting of position 71, position 95, position 101 and position 102.
6. The nucleic acid of claim 5, wherein the mutated Tn10-derived Tet repressor comprises the amino acid sequence corresponding to positions 1 to 207 of SEQ ID NO: 2.
7. The nucleic acid of claim 1, wherein the second polypeptide comprises a transcription activation domain of herpes simplex virion protein 16.
8. The nucleic acid of claim 7, wherein the transcription activation domain comprises a C-terminal amino acid region of herpes simplex virus virion protein 16 comprising an amino acid sequence corresponding to positions 208 to 335 of SEQ ID NO: 2.
9. The nucleic acid of claim 7, wherein the transcription activation domain comprises at least one copy of a C-terminal region of herpes simplex virus virion protein 16 comprising an amino acid sequence shown in SEQ ID NO: 4.
10. The nucleic acid of claim 1, wherein the second polypeptide comprises a transcription activation domain selected from the group consisting of: an acidic transcription activation domain, a proline-rich transcription activation domain, a serine/threonine-rich transcription activation domain and a glutamine-rich transcription activation domain.

11. The nucleic acid molecule of claim 1, wherein the fusion protein further comprises an operatively linked third polypeptide which promotes transport of the fusion protein to a cell nucleus.
- 5 12. A fusion protein which activates transcription comprising a first polypeptide which binds to a *tet* operator sequence in the presence of tetracycline or a tetracycline analogue operatively linked to a second polypeptide which activates transcription in eukaryotic cells.
- 10 13. The fusion protein of claim 12, wherein the first polypeptide is a mutated Tet repressor.
14. The fusion protein of claim 12, wherein the second polypeptide comprises a transcription activation domain of herpes simplex virion protein 16.
- 15 15. The fusion protein of claim 12, which comprises an amino acid sequence shown in SEQ ID NO: 2.
16. A recombinant vector comprising the nucleic acid molecule of claim 1 in a form suitable for expression of the fusion protein in a host cell.
- 20 17. The recombinant vector of claim 16, wherein expression of the fusion protein is regulated by at least one tissue-specific regulatory element.
18. The recombinant vector of claim 16, wherein expression of the fusion protein is regulated by at least one *tet* operator sequence.
- 25 19. A host cell comprising the recombinant vector of claim 16.
20. The host cell of claim 19, further comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.
- 30 21. The host cell of claim 20, wherein the nucleotide sequence to be transcribed is an exogenous nucleotide sequence introduced into the host cell.
- 35 22. The host cell of claim 20, wherein the nucleotide sequence to be transcribed is an endogenous nucleotide sequence to which at least one *tet* operator sequence has been operatively linked.
23. The host cell of claim 20, which is a mammalian cell.

24. The host cell of claim 21, which is a human cell.
25. The host cell of claim 20, which is a yeast, insect or fungal cell.
- 5 26. A method for stimulating transcription of the nucleotide sequence operatively linked to the at least one *tet* operator sequence in the host cell of claim 20 comprising contacting the host cell with tetracycline or a tetracycline analogue.
- 10 27. The method of claim 26, wherein the tetracycline analogue is anhydrotetracycline, doxycycline or cyanotetracycline.
28. The method of claim 26, further comprising isolating a protein encoded by the nucleotide sequence operatively linked to the at least one *tet* operator sequence from the host
15 cells or from a culture medium in which the host cells are grown.
29. A non-human transgenic organism comprising a transgene comprising the nucleic acid molecule of claim 1 in a form suitable for expression of the fusion protein in cells of the non-human transgenic organism.
- 20 30. The non-human transgenic organism of claim 29, wherein the transgene is integrated at a predetermined location within the genome of the organism.
31. The non-human transgenic organism of claim 29, further comprising a nucleotide
25 sequence to be transcribed operatively linked to at least one *tet* operator sequence.
32. The non-human transgenic organism of claim 30, further comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.
- 30 33. A method for stimulating transcription of the nucleotide sequence operatively linked to the at least one *tet* operator sequence in the non-human transgenic organism of claim 31, comprising administering tetracycline or a tetracycline analogue to the organism.
34. The method of claim 33, wherein the tetracycline analogue is anhydrotetracycline,
35 doxycycline or cyanotetracycline.
35. An isolated nucleic acid encoding a fusion protein which inhibits transcription in eukaryotic cells, the fusion protein comprising a first polypeptide which binds to a *tet*

operator sequence operatively linked to a heterologous second polypeptide which inhibits transcription in eukaryotic cells.

5 36. The nucleic acid of claim 35, wherein the first polypeptide binds to a *tet* operator sequence in the absence but not the presence of tetracycline or a tetracycline analogue.

37. The nucleic acid of claim 36, wherein the first polypeptide is a Tet repressor.

10 38. The nucleic acid of claim 37, wherein the first polypeptide comprises an amino acid sequence shown in SEQ ID NO: 17.

39. The nucleic acid of claim 35, wherein the first polypeptide binds to a *tet* operator sequence in the presence but not the absence of tetracycline or a tetracycline analogue.

15 40. The nucleic acid of claim 39, wherein the first polypeptide is a mutated Tet repressor.

41. The nucleic acid of claim 40, wherein the mutated Tet repressor has at least one amino acid substitution compared to a wild-type Tet repressor.

20 42. The nucleic acid of claim 40, wherein the mutated Tet repressor has at least one amino acid addition or deletion compared to a wild-type Tet repressor.

25 43. The nucleic acid of claim 41, wherein the mutated Tet repressor has an amino acid substitution at at least one amino acid position corresponding to an amino acid position selected from the group consisting of position 71, position 95, position 101 and position 102 of a wild-type Tn10-derived Tet repressor amino acid sequence.

30 44. The nucleic acid of claim 43, wherein the mutated Tet repressor comprises an amino acid sequence shown in SEQ ID NO: 19.

45. The nucleic acid of claim 35, wherein the second polypeptide comprises a transcription silencer domain of a v-erbA oncogene product.

35 46. The nucleic acid of claim 45, wherein the second polypeptide comprises an amino acid sequence shown in SEQ ID NO: 23.

47. The nucleic acid of claim 35, wherein the second polypeptide comprises a transcription silencer domain of a Drosophila Krueppel protein.

48. The nucleic acid of claim 47, wherein the second polypeptide comprises an amino acid sequence shown in SEQ ID NO: 21.
49. The nucleic acid of claim 35, wherein the second polypeptide comprises a transcription silencer domain of a protein selected from the group consisting of the retinoic acid receptor alpha, the thyroid hormone receptor alpha, the yeast Ssn6/Tup1 protein complex, the Drosophila protein even-skipped, SIR1, NeP1, the Drosophila dorsal protein, TSF3, SFI, the Drosophila hunchback protein, the Drosophila knirps protein, WT1, Oct-2.1, the Drosophila engrailed protein, E4BP4 and ZF5.
50. The nucleic acid molecule of claim 35, wherein the fusion protein further comprises an operatively linked third polypeptide which promotes transport of the fusion protein to a cell nucleus.
51. A fusion protein which inhibits transcription in eukaryotic cells, comprising a first polypeptide which binds to a *tet* operator sequence operatively linked to a heterologous second polypeptide which inhibits transcription in eukaryotic cells.
52. The fusion protein of claim 51, wherein the first polypeptide binds to a *tet* operator sequence in the absence but not the presence of tetracycline or a tetracycline analogue.
53. The fusion protein of claim 52, wherein the first polypeptide comprises an amino acid sequence shown in SEQ ID NO: 17.
54. The fusion protein of claim 51, wherein the first polypeptide binds to a *tet* operator sequence in the presence but not the absence of tetracycline or a tetracycline analogue.
55. The fusion protein of claim 54, wherein the first polypeptide comprises an amino acid sequence shown in SEQ ID NO: 19.
56. The fusion protein of claim 51, wherein the second polypeptide comprises a transcription silencer domain of a v-erbA oncogene product.
57. The fusion protein of claim 56, wherein the second polypeptide comprises an amino acid sequence shown in SEQ ID NO: 23.
58. The fusion protein of claim 51, wherein the second polypeptide comprises a transcription silencer domain of a Drosophila Krueppel protein.

59. The fusion protein of claim 58, wherein the second polypeptide comprises an amino acid sequence shown in SEQ ID NO: 21.

60. The fusion protein of claim 51, wherein the second polypeptide comprises a transcription silencer domain of a protein selected from the group consisting of the retinoic acid receptor alpha, the thyroid hormone receptor alpha, the yeast Ssn6/Tup1 protein complex, the Drosophila protein even-skipped, SIR1, NeP1, the Drosophila dorsal protein, TSF3, SFI, the Drosophila hunchback protein, the Drosophila knirps protein, WT1, Oct-2.1, the Drosophila engrailed protein, E4BP4 and ZF5.

61. A recombinant vector comprising the nucleic acid molecule of claim 35 in a form suitable for expression of the fusion protein in a host cell.

62. A host cell comprising the recombinant vector of claim 61.

63. The host cell of claim 62, further comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.

64. The host cell of claim 63, wherein the nucleotide sequence to be transcribed is an exogenous nucleotide sequence introduced into the host cell.

65. The host cell of claim 63, wherein the nucleotide sequence to be transcribed is an endogenous nucleotide sequence to which at least one *tet* operator sequence has been operatively linked.

66. A method for modulating transcription of the nucleotide sequence operatively linked to the at least one *tet* operator sequence in the host cell of claim 63, comprising modulating the concentration of tetracycline or a tetracycline analogue that is in contact with the host cell.

67. A non-human transgenic organism comprising a transgene comprising the nucleic acid molecule of claim 35 in a form suitable for expression of the fusion protein in cells of the non-human transgenic organism.

68. The non-human transgenic organism of claim 67, wherein the transgene is integrated at a predetermined location within the genome of the organism.

69. The non-human transgenic organism of claim 67, further comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.

70. The non-human transgenic organism of claim 68, further comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.
- 5 71. A method for modulating transcription of the nucleotide sequence operatively linked to the at least one *tet* operator sequence in the non-human transgenic organism of claim 69, comprising modulating the concentration of tetracycline or a tetracycline analogue present in the organism.
- 10 72. The method of claim 71, wherein the tetracycline analogue is anhydrotetracycline, doxycycline or cyanotetracycline.
73. A host cell comprising:
a first nucleic acid encoding a first fusion protein which activates
15 transcription, the first fusion protein comprising a first polypeptide which binds to a *tet* operator sequence operatively linked to a second polypeptide which activates transcription in eukaryotic cells;
a second nucleic acid encoding a second fusion protein which inhibits
transcription, the second fusion protein comprising a third polypeptide which binds to a *tet*
20 operator sequence operatively linked to a fourth polypeptide which inhibits transcription in eukaryotic cells; and
a third nucleic acid molecule comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.
- 25 74. The host cell of claim 73, wherein the first polypeptide binds to a *tet* operator sequence in the presence, but not the absence of tetracycline or a tetracycline analogue and the third polypeptide binds to a *tet* operator sequence in the absence, but not the presence of tetracycline or a tetracycline analogue.
- 30 75. The host cell of claim 73, wherein the first polypeptide binds to a *tet* operator sequence in the absence, but not the presence of tetracycline or a tetracycline analogue and the third polypeptide binds to a *tet* operator sequence in the presence, but not the absence, of tetracycline or a tetracycline analogue.
- 35 76. A non-human transgenic organisms comprising:
a first transgene encoding a first fusion protein which activates transcription, the first fusion protein comprising a first polypeptide which binds to a *tet* operator sequence operatively linked to a second polypeptide which activates transcription in eukaryotic cells;

a second transgene encoding a second fusion protein which inhibits transcription, the second fusion protein comprising a third polypeptide which binds to a *tet* operator sequence operatively linked to a fourth polypeptide which inhibits transcription in eukaryotic cells; and

- 5 a third transgene comprising a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.

77. The non-human transgenic animal of claim 76, wherein the first polypeptide binds to a *tet* operator sequence in the presence, but not the absence of tetracycline or a tetracycline
10 analogue and the third polypeptide binds to a *tet* operator sequence in the absence, but not the presence of tetracycline or a tetracycline analogue.

78. The non-human transgenic animal of claim 76, wherein the first polypeptide binds to a *tet* operator sequence in the absence, but not the presence of tetracycline or a tetracycline
15 analogue and the third polypeptide binds to a *tet* operator sequence in the presence, but not the absence, of tetracycline or a tetracycline analogue.

79. A recombinant vector for coordinate, bidirectional transcription of a first and a second nucleotide sequence to be transcribed, the vector comprising a nucleotide sequence
20 comprising in a 5' to 3' direction:

- a first cloning site for introduction of a first nucleotide sequence to be transcribed, which is operatively linked to
 at least one *tet* operator sequence, which is operatively linked to
25 a second cloning site for introduction of a second nucleotide sequence to be transcribed,

wherein transcription of the first and second nucleotide sequence introduced into the vector proceeds in opposite directions relative to the at least one *tet* operator sequence.

30

80. The recombinant vector of claim 79, wherein the first and second nucleotide sequence to be transcribed have been introduced into the first and second cloning sites, respectively.

81. The recombinant vector of claim 80, wherein the first and second nucleotide
35 sequences encodes polypeptide subunits of a heterodimeric protein.

82. The recombinant vector of claim 80, wherein the first nucleotide sequence encodes a protein of interest and the second nucleotide sequence encodes a detectable marker.

83. At least one recombinant vector for independent regulation of transcription of a first and a second nucleotide sequence to be transcribed, the at least one vector comprising a nucleotide sequence comprising:

- 5 a first cloning site for introduction of a first nucleotide sequence to be transcribed, operatively linked to at least one *tet* operator sequence of a first class type; and
a second cloning site for introduction of a second nucleotide sequence to be transcribed, operatively linked to at least one *tet* operator sequence of a second class type.

10 84. The at least one recombinant vector of claim 83, wherein either the first or the second class type of *tet* operator sequence is a class B *tet* operator sequence.

15 85. The at least one recombinant vector of claim 83, wherein the first and second nucleotide sequence to be transcribed have been introduced into the first and second cloning site, respectively.

86. The at least one recombinant vector of claim 85, wherein the first nucleotide sequence comprises a therapeutic gene and the second nucleotide sequence comprises a suicide gene.

20 87. A kit comprising a carrier means having in close confinement therein at least two container means comprising:


- a first container means containing a first nucleic acid encoding a fusion protein which activates transcription, the fusion protein comprising a first polypeptide which binds to a *tet* operator sequence in the presence of tetracycline or a tetracycline analogue operatively linked to a second polypeptide which activates transcription in eukaryotic cells; and
25 a second container means containing a second nucleic acid comprising a cloning site for introduction of a nucleotide sequence to be transcribed operatively linked to at least one *tet* operator sequence.

30 88. The kit of claim 87, further comprising a third container means containing a third nucleic acid encoding a fusion protein which inhibits transcription, the fusion protein comprising a first polypeptide which binds to a *tet* operator sequence operatively linked to a heterologous second polypeptide which inhibits transcription in eukaryotic cells.

35 89. The kit of claim 88, further comprising a fourth container means containing a tetracycline or a tetracycline analogue.

1 / 13


FIG. 1



SUBSTITUTE SHEET (RULE 26)


2 / 13

FIG. 2



SUBSTITUTE SHEET (RULE 26)


FIG. 3



SUBSTITUTE SHEET (RULE 26)

4 / 13

FIG. 4A




SUBSTITUTE SHEET (RULE 26)

5.6.1

| | A1 | A2 | B1 | B2 | C1 | C2 | D1 | D2 | E1 | E2 | |
|----|----|----|----|----|-----|----|----|----|----|----|---|
| 1 | A | A | A | T | T | A | A | C | T | T | T |
| 2 | C | C | T | T | A | T | C | A | G | T | G |
| 3 | A | A | A | A | C | A | A | A | A | A | A |
| 4 | C | A | T | T | A | T | C | A | T | A | A |
| 5 | T | T | G | A | A | T | A | G | T | C | A |
| 6 | G | A | C | T | A | T | T | G | T | T | C |
| 7 | T | T | G | A | A | T | A | G | T | C | A |
| 8 | T | T | G | A | A | T | A | G | T | C | A |
| 9 | T | T | G | A | A | T | A | G | T | C | A |
| 10 | T | T | G | A | A | T | A | G | T | C | A |
| 11 | T | T | G | A | A | T | A | G | T | C | A |
| 12 | T | T | G | A | A | T | A | G | T | C | A |
| 13 | T | T | G | A | A | T | A | G | T | C | A |
| 14 | T | T | G | A | A | T | A | G | T | C | A |
| 15 | T | T | G | A | A | T | A | G | T | C | A |
| 16 | T | T | G | A | A | T | A | G | T | C | A |
| 17 | T | T | G | A | A | T | A | G | T | C | A |
| 18 | T | T | G | A | A | T | A | G | T | C | A |
| 19 | T | T | G | A | A | T | A | G | T | C | A |
| 20 | T | T | G | A | A | T | A | G | T | C | A |
| 21 | T | T | G | A | A | T | A | G | T | C | A |
| 22 | T | T | G | A | A | T | A | G | T | C | A |
| 23 | T | T | G | A | A | T | A | G | T | C | A |
| 24 | T | T | G | A | A | T | A | G | T | C | A |
| 25 | T | T | G | A | A | T | A | G | T | C | A |
| 26 | T | T | G | A | A | T | A | G | T | C | A |
| 27 | T | T | G | A | A | T | A | G | T | C | A |
| 28 | T | T | G | A | A | T | A | G | T | C | A |
| 29 | T | T | G | A | A | T | A | G | T | C | A |
| 30 | T | T | G | A | A | T | A | G | T | C | A |
| 31 | T | T | G | A | A | T | A | G | T | C | A |
| 32 | T | T | G | A | A | T | A | G | T | C | A |
| 33 | T | T | G | A | A | T | A | G | T | C | A |
| 34 | T | T | G | A | A | T | A | G | T | C | A |
| 35 | T | T | G | A | A | T | A | G | T | C | A |
| 36 | T | T | G | A | A | T | A | G | T | C | A |
| 37 | T | T | G | A | A | T | A | G | T | C | A |
| 38 | T | T | G | A | A | T | A | G | T | C | A |
| 39 | T | T | G | A | A | T | A | G | T | C | A |
| 40 | T | T | G | A | A | T | A | G | T | C | A |
| 41 | T | T | G | A | A | T | A | G | T | C | A |
| 42 | T | T | G | A | A | T | A | G | T | C | A |
| 43 | T | T | G | A | A | T | A | G | T | C | A |
| 44 | T | T | G | A | A | T | A | G | T | C | A |
| 45 | T | T | G | A | A | T | A | G | T | C | A |
| 46 | T | T | G | A | A | T | A | G | T | C | A |
| 47 | T | T | G | A | A | T | A | G | T | C | A |
| 48 | T | T | G | A | A | T | A | G | T | C | A |
| 49 | T | T | G | A | A | T | A | G | T | C | A |
| 50 | T | T | G | A | A</ | | | | | | |

6 / 13

FIG. 6



SUBSTITUTE SHEET (RULE 26)

7 / 13

FIG. 7A

5' GAATTCGGGG
EcoRI + 75

CCGCGGAGGCTGGATCGGTCCCGGTGTCTTCTATGGAGGTCAAACAGCGTGGA

+ 1
←
C

P_{hcmv}*-3TGGCGTCTCCAGGCGATCTGACGGTTCCTAAACGAGCTCTGCTTATATAGG
-31*tet O*TC (GAGTTTACCACTCCCTATCAGTGATAGAGAAAAGTGAAAGTC)₇GAGCP_{hcmv}*-1TCGGTACCCGGGTCGAGTAGGCGTGTACGGTGGGAGGCCTATATAAGCAGAG
-53CTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGA
+ 1 →CCTCCATAGAAGACACCGGGACCGATCCAGCCTCCGCGGCCCGAATTC 3'
+ 75 EcoRI

SUBSTITUTE SHEET (RULE 26)

9 / 13

FIG. 8A





FIG. 8B



SUBSTITUTE SHEET (RULE 26)

FIG. 9A




FIG. 9B






FIG. 10




SUBSTITUTE SHEET (RULE 26)

FIG. 11



SUBSTITUTE SHEET (RULE 26)

FIG. 12



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 95/08179

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/00 C12N15/12 C12N15/62 C12N15/63 C12N15/67
C12N15/85 C12N5/10 C12N1/15 C12N1/19 C07K14/245
A01K61/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|------------------------------|
| Y | TRENDS IN BIOCHEMICAL SCIENCES, vol. 19, no. 1, January 1994 ELSEVIER, AMSTERDAM, NL, pages 38-42, I.G. COWELL 'Repression versus activation in the control of gene transcription' see the whole document --- | 35-38, 51-53, 61-64,66 |
| Y | TRENDS IN BIOCHEMICAL SCIENCES, vol. 18, no. 12, December 1993 ELSEVIER, AMSTERDAM, NL, pages 471-475, M. GOSSEN ET AL. 'Control of gene activity in higher eucaryotic cells by procaryotic regulatory elements' see the whole document --- -/-- | 35-38, 51-53, 61-64,66 |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

3 November 1995

Date of mailing of the international search report

24. 11. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Hornig, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/08179

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|------------------------------|
| Y | <p>PROC. NATL. ACAD. SCI. U. S. A. (1992), 89(12), 5547-51 CODEN: PNASA6;ISSN: 0027-8424, 15 June 1992 GOSSEN, MANFRED ET AL 'Tight control of gene expression in mammalian cells by tetracycline -responsive promoters' cited in the application the whole document</p> <p>---</p> | 35-38, 51-53, 61-64,66 |
| A | <p>WO,A,94 04672 (DNX CORP ;BYRNE GUERARD (US)) 3 March 1994 cited in the application see the whole document</p> <p>---</p> | 1-89 |
| A | <p>J. BACTERIOL., vol. 175, no. 4, February 1993 AM. SOC. MICROBIOL.,BALTIMORE,US;, pages 1206-1210, B. HECHT ET AL. 'Noninducible Tet repressor mutations map from the operator binding motif to the C terminus' cited in the application see the whole document</p> <p>---</p> | 1-89 |
| A | <p>THE PLANT JOURNAL, vol. 2, no. 3, May 1992 BLACKWELL, OXFORD, UK, pages 397-404, C. GATZ ET AL. 'Stringent repression and homogeneous de-repression by tetracycline of a modified CaMV 35S promoter in intact transgenic tobacco plants' cited in the application see the whole document</p> <p>---</p> | 1-89 |
| A | <p>PLANT J. (1994), 5(4), 559-69 CODEN: PLJUED;ISSN: 0960-7412, 1994 WEINMANN, PAMELA ET AL 'a chimeric transactivator allows tetracycline -responsive gene expression in whole plants' see the whole document</p> <p>---</p> | 1-89 |
| P,A | <p>WO,A,94 29442 (BASF AG ;BUJARD HERMANN (DE); GOSSEN MANFRED (DE); SALFELD JOCHEN) 22 December 1994 see the whole document</p> <p>---</p> | 1-89 |
| | <p>---</p> <p>-/--</p> | |

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/08179

| C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|--|---|----------------------------------|
| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| P,X | <p>SCIENCE (WASHINGTON, D. C.) (1995), 268(5218), 1766-9 CODEN: SCIEAS;ISSN: 0036-8075, 23 June 1995 GOSSEN, MANFRED ET AL 'Transcriptional activation by tetracyclines in mammalian cells' the whole document</p> <p style="text-align: center;">---</p> | 1-28 |
| P,X | <p>KEYSTONE SYMPOSIUM ON GENE THERAPY AND MOLECULAR MEDICINE, STEAMBOAT SPRINGS, COLORADO, USA, MARCH 26-APRIL 1, 1995. JOURNAL OF CELLULAR BIOCHEMISTRY SUPPLEMENT 0 (21A). 1995. 379. ISSN: 0733-1959, ABSTRACT NO. C6-220, LIANG X ET AL 'Enhanced and switchable expression systems for gene-transfer.' see abstract</p> <p style="text-align: center;">---</p> | 83-86 |
| P,X | <p>KEYSTONE SYMPOSIUM ON GENE THERAPY AND MOLECULAR MEDICINE, STEAMBOAT SPRINGS, COLORADO, USA, MARCH 26-APRIL 1, 1995. JOURNAL OF CELLULAR BIOCHEMISTRY SUPPLEMENT 0 (21A). 1995. 355. ISSN: 0733-1959, GOSSEN M ET AL 'Exploiting prokaryotic elements for the control of gene activity in higher eukaryotics. March 28, 1995;' abstract no. C6-004 see abstract</p> <p style="text-align: center;">---</p> | 1-4,7, 10, 12-14, 16-28 |
| P,X | <p>MOL. CELL. BIOL. (1995), 15(4), 1907-14 CODEN: MCEBD4;ISSN: 0270-7306, April 1995 DEUSCHLE, ULRICH ET AL 'Tetracycline -reversible silencing of eukaryotic promoters' see the whole document</p> <p style="text-align: center;">-----</p> | 35-38, 51-53, 61-64,66 |

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/US 95/08179

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|---|----------------------------------|
| WO-A-9404672 | 03-03-94 | AU-B- 5099393 CA-A- 2143326 EP-A- 0665883 | 15-03-94 03-03-94 09-08-95 |
| WO-A-9429442 | 22-12-94 | AU-B- 7108194 | 03-01-95 |


Form PCT/ISA/210 (patent family annex) (July 1992)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|--|---|
| (51) International Patent Classification ⁶ : C12N 15/00, 15/12, 15/62, 15/63, 15/67, 15/85, 5/10, 1/15, 1/19, C07K 14/245, A01K 61/027 | | A1 | (11) International Publication Number: WO 96/01313 |
| | | | (43) International Publication Date: 18 January 1996 (18.01.96) |
| (21) International Application Number: PCT/US95/08179 | | Manfred [DE/US]; 978 Arlington Boulevard #B, El Cerrito, CA 94530 (US). | |
| (22) International Filing Date: 29 June 1995 (29.06.95) | | (74) Agents: DECONTI, Giulio, A., Jr. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US). | |
| (30) Priority Data: | | (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). | |
| 08/270,637 1 July 1994 (01.07.94) US 08/275,876 15 July 1994 (15.07.94) US 08/383,754 3 February 1995 (03.02.95) US 08/486,814 7 June 1995 (07.06.95) US | | | |
| (60) Parent Applications or Grants | | Published | |
| (63) Related by Continuation | | With international search report. | |
| US 08/270,637 (CIP) | | Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. | |
| Filed on 1 July 1994 (01.07.94) | | | |
| US 08/275,876 (CIP) | | | |
| Filed on 15 July 1994 (15.07.94) | | | |
| US 08/383,754 (CIP) | | | |
| Filed on 3 February 1995 (03.02.95) | | | |
| US 08/486,814 (CIP) | | | |
| Filed on 7 June 1995 (07.06.95) | | | |
| (71)(72) Applicants and Inventors: BUIARD, Hermann [DE/DE]; Remler Strasse 9, D-69120 Heidelberg (DE). GOSSEN, | | | |

(54) Title: TETRACYCLINE-REGULATED TRANSCRIPTIONAL MODULATORS



(57) Abstract

Nucleic acid molecules and proteins useful for regulating the expression of genes in eukaryotic cells and organisms are disclosed. The invention provides a transcriptional activator fusion protein which binds to *tet* operator sequences and stimulates transcription of a *tet* operator-linked gene in the presence, but not the absence, of tetracycline (or analogue thereof). The invention further provides transcriptional inhibitor fusion proteins which inhibit transcription of a *tet* operator-linked gene in a regulated manner. In one embodiment, the inhibitor fusion protein binds to *tet* operator sequences in the absence, but not the presence, of tetracycline (or analogue). In another embodiment, the inhibitor fusion protein binds to *tet* operator sequences in the presence, but not the absence, of tetracycline (or analogue). The transcriptional activator and inhibitor fusion proteins of the invention can be used in combination to regulate expression of one or multiple *tet* operator-linked genes. Novel *tet* operator-containing transcription units which allow for coordinate or independent tetracycline-regulated expression of two or more genes by the transcriptional modulators of the invention are also disclosed. Kits including the components of the regulatory system are also encompassed by the invention.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria | GB | United Kingdom | MR | Mauritania |
| AU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | IT | Italy | PL | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgyzstan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SI | Slovenia |
| CI | Côte d'Ivoire | LI | Liechtenstein | SK | Slovakia |
| CM | Cameroon | LK | Sri Lanka | SN | Senegal |
| CN | China | LU | Luxembourg | TD | Chad |
| CS | Czechoslovakia | LV | Latvia | TG | Togo |
| CZ | Czech Republic | MC | Monaco | TJ | Tajikistan |
| DE | Germany | MD | Republic of Moldova | TT | Trinidad and Tobago |
| DK | Denmark | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | US | United States of America |
| FI | Finland | MN | Mongolia | UZ | Uzbekistan |
| FR | France | | | VN | Viet Nam |
| GA | Gabon | | | | |